

Paleontological Research



Papers in Press

“Papers in Press” includes peer-reviewed, accepted manuscripts of research articles, reviews, and short notes to be published in *Paleontological Research*. They have not yet been copy edited and/or formatted in the publication style of *Paleontological Research*. As soon as they are printed, they will be removed from this website. Please note they can be cited using the year of online publication and the DOI, as follows:

Humblet, M. and Iryu, Y. 2014: Pleistocene coral assemblages on Irabu-jima, South Ryukyu Islands, Japan. *Paleontological Research*, doi: 10.2517/2014PR020.

The Guadalupian (Permian) Gufeng Formation on the north margin of the South China block: A review of the lithostratigraphy, radiolarian biostratigraphy, and geochemical characteristics

TSUYOSHI ITO¹, KOJI U. TAKAHASHI², ATSUSHI MATSUOKA³ AND QINGLAI FENG⁴

¹*Research Institute of Geology and Geoinformation, Geological Survey of Japan, AIST, Tsukuba, Ibaraki 305-8567, Japan (e-mail: ito-t@aist.go.jp)*

²*Research Institute of Geo-Resources and Environment, Geological Survey of Japan, AIST, Tsukuba, Ibaraki 305-8567, Japan*

³*Department of Geology, Faculty of Science, Niigata University, Niigata 950-2181, Japan*

⁴*State Key Laboratory of Geo-processes and Mineral Resources, China University of Geosciences, Wuhan 430074, China*

Abstract.

The Guadalupian (Permian) Gufeng Formation, distributed over the north margin of the South China block, is characterized by siliceous rock with muddy rock and yields several fossils representative by radiolarian fossils. The lithostratigraphy, radiolarian occurrences with biostratigraphy, and geochemical characteristics of the Gufeng Formation are reviewed and summarized in this paper. The distributional area of the Gufeng Formation is subdivided into the Lower, Middle, and Upper Yangtze regions. The Gufeng Formation in the Lower Yangtze region is lithostratigraphically characterized by muddy rock, chert, and muddy rock with nodules in descending order. The Gufeng Formation in the Middle Yangtze region is lithostratigraphically characterized by chert, muddy rock with carbonate, and muddy rock in ascending order. The lithostratigraphic characteristics of the “Gufeng Formation” in the south margin greatly differ from the typical Gufeng Formation in containing fewer siliceous rocks and much clastic rocks and in having a significantly thicker total thickness; therefore, this article considers that the “Gufeng Formation” in the south margin should be isolated from the Gufeng Formation in the north margin. The radiolarian biostratigraphy of the Lower and Middle Yangtze regions is composed of the *Pseudoalbaillella*

globosa, *Ps. monacantha*, and *Follicucullus scholasticus-Ruzhencevispongus uralicus* assemblage zones in ascending order. The lower Gufeng Formation has abundantly yielded the *Albaillellaria*; however, its occurrence ratio decreases stratigraphically upward. The geochemical characteristics indicate the following facts: (1) the siliceous rocks are generally biogenic; (2) even though a previous study concluded that the organic matter within the cherts was originated from terrestrial or reworked organic matter, the origin remains debatable; and (3) the Gufeng Formation was formed on a continental shelf. The geochemical characteristics, in addition to the lithological characteristics, indicate that the redox conditions of the Gufeng Formation in the Lower Yangtze region changed from aerobic to suboxic to anoxic, while that in the Middle Yangtze region changed from oxic to anoxic to suboxic.

Key words: Gufeng Formation, lithology, radiolarian biostratigraphy, sedimentary environment, South China block

Introduction

The Permian period is one of the most varying intervals during Earth's history (e.g. Hein, 2004) because several events occurred during this period, such as the most severe mass extinction in the Phanerozoic at the Guadalupian-Lopingian boundary and the end-Permian (Jin *et al.*, 1994; Stanley and Yang, 1994), multiple glaciations (e.g. Isbell *et al.*, 2011), a global cooling event called as the Kamura event (e.g. Isozaki *et al.*, 2007a, b), and a large amount of chert accumulation in the middle Permian called as the mid-Permian chert event (Murchey and Jones, 1992) or Permian Chert Event (Beauchamp and Baud, 2002).

The Gufeng (Kuhfeng) Formation in South China has yielded Guadalupian (middle Permian) fossils in several taxa (e.g. ammonoids, brachiopods, and conodonts). In particular, the Gufeng Formation is known to have abundant radiolarian-bearing strata. This fossiliferous formation has attracted attention in recent years (e.g. Zhang *et al.*, in press) because the formation may record paleobiological responses to environmental changes through the Permian. Yao *et al.* (2015) studied fluctuation of chert bed thickness of the Gufeng Formation and performed astronomical cyclostratigraphic study with radiolarian productivities.

In addition, the Gufeng Formation is important from paleogeographic and paleobiogeographic points of view. The Gufeng Formation consists of sediments on the South China block (Yangtze block). This block had been situated at the east end of the Paleotethys near the boundary between the superoceans Panthalassa and Paleotethys in the middle Permian (e.g. Ziegler *et al.*, 1997; Golonka and Ford, 2000) (Figure 1). Because the South China block acted as a wall between the both superoceans, a paleobiogeographic boundary has been recognized around the South China block in several taxa, such as fusulinid (e.g. Kobayashi, 1997), brachiopod (e.g. Shen *et al.*, 2009), and radiolaria (Ito *et al.*, 2016b).

Furthermore, the Gufeng Formation is known to have oil-rich strata; therefore, petrochemical studies have been performed (e.g. Xia *et al.*, 1995; Zeng *et al.*, 2004; Yang and Yao, 2008). Previous studies have focused on the porosity of the rocks in the Gufeng Formation as a reservoir rock (Wu *et al.*, 2012; Liang *et al.*, 2014). In recent years, its potential as an origin and reservoir of shale gas has also been a focus of study (Du *et al.*, 2015).

In short, the strata of the Gufeng Formation are extremely significant in terms of paleontology, sedimentology, paleogeography, paleobiogeography, and petroleum geology. Meanwhile, the formation has regional differences in several characteristics,

such as lithostratigraphy and biostratigraphy. Due to these regional differences, even the definition and distributions of the formation have been debated (e.g. Hu, 2000; Kuwahara *et al.*, 2004; Yao *et al.*, 2007). On the basis of previous studies and combined with our study, here we compile the lithostratigraphy, radiolarian occurrences with biostratigraphy, and geochemical characteristics of the Gufeng Formation in each region. This article aims to provide a general review of these aspects for future reference.

Lithostratigraphy

Definition and distribution

The Gufeng Formation was originally called as the Gufeng Zhen Limestone at the Hujia Village of Gufeng Town, Anhui Province by Ye and Li (1924) (Figure 2). Bureau of Geology and Mineral Resources of Anhui Province (1997) redefined the Gufeng Formation as follows: “It is characterized by black or gray-yellow thin-bedded siliceous rocks, siliceous shales, siltstones, carbonous shales, and manganese shales. The Gufeng Formation overlies the Qixia Formation and underlies the Longtan Formation. Basal shales include manganese and phosphate nodules. Several taxa of fossils (e.g. ammonoids, brachiopods, bivalves, and radiolarians) occur from it.”

The bureaus of geology and mineral resources in each province in China have compiled the regional geology in each province. The columnar sections of the stratotype and hypostratotype sections established by the bureaus with some notable sections are summarized in Figure 3 and Figure 4. Bureau of Geology and Mineral Resources of Anhui Province (1997) established the Hujia section (30°50'E, 118°24'E) in Gufeng as a stratotype section of the Gufeng Formation. In other provinces, the bureaus of geology and mineral resources in each province established hypostratotype sections of the Gufeng Formation: the Lengwu section (29°55'E, 119°32'E) in Tonglu (Bureau of Geology and Mineral Resources of Zhejiang Province, 1996), the Hechong section (32°09'E, 119°06'E) in Jurong (Bureau of Geology and Mineral Resources of Jiangsu Province, 1997), the Bijiashan section (30°12'E, 114°47'E) in Daye (Bureau of Geology and Mineral Resources of Hubei Province, 1996), and the Lengshuixi section in Shizhu (Bureau of Geology and Mineral Resources of Sichuan Province, 1997).

The Gufeng Formation is distributed over the north margin of the South China block (Figure 3). Its distributional area is subdivided into three regions: the Lower Yangtze region (north Zhejiang, south Jiangsu, and south Anhui), the Middle Yangtze region (Hubei), and the Upper Yangtze region (Shaanxi and north Sichuan). Previous studies have also considered siliceous rocks on the south margin of the South China block

(south Hunan, northwest Guangdong, and east Guangxi) to be part of the Gufeng Formation. However, the so-called “Gufeng Formation” in the south margin differs from the typical Gufeng Formation in the north margin in lithostratigraphy, as will be described in detail. We therefore regard the Gufeng Formation as being distributed only over the north margin.

Lithology and its correlations

North margin.—The Hujia section in Gufeng, the stratotype section of the Gufeng Formation, is divided into the following subsections in ascending order (Bureau of Geology and Mineral Resources of Anhui Province, 1997): subsection 1: brownish-black manganese mudstone; subsection 2: yellow mudstone, interbedding gray siliceous mudstone and thin-bedded chert; subsection 3: yellow mudstone including gray phosphate muddy nodules; subsection 4: gray siliceous mudstone; subsection 5: black thin-bedded chert interbedding black thin-bedded siliceous mudstone; and subsection 6: black thin-bedded siliceous mudstone interbedding carbonaceous mudstone. The Gufeng Formation of the section conformably contacts with the underlying Qixia and overlying Longtan formations.

The Gufeng Formation at Chaohu in Anhui has been studied in detail mainly by a

research group of Nagoya University and Nanjing University (Nagai and Zhu, 1992; Nagai *et al.*, 1998; Zhu *et al.*, 1999; Kametaka *et al.*, 2002, 2005, 2009; Takebe *et al.*, 2007). Kametaka *et al.* (2002) and Kametaka *et al.* (2005) divided the Gufeng Formation of the Anmenkou section at Chaohu into two members: the Phosphate Nodule-bearing Mudstone Member (PNMM) and the Siliceous Rock Member (SRM) in ascending order. The former (*ca.* 4 m in total thickness) consists of mudstone including abundant phosphate nodules; the latter (*ca.* 25 m) is composed primarily of alternations of black chert, mudstone, and siliceous mudstone, with minor tuffaceous mudstone and porous chert beds. Ito *et al.* (2013a) investigated the Liuhuang section near the Anmenkou section and presented lithologic similarity between the Liuhuang and Anmenkou sections. In the Liuhuang section, the siliceous rock is dark-colored (black to dark-gray) and well bedded (Figure 5A, B). The Gufeng Formation disconformably contacts with the limestone of the underlying Maokou Formation in the Liuhuang section (Figure 5C). Zhang *et al.* (in press) investigated the Hexian core drilled at Hexian 50 km northeast of Chaohu and subdivided the Gufeng Formation in the core into three members, the Lower Phosphate Nodule-bearing Mudstone Member (LPMM), the Lower Chert-Mudstone Member (LCMM), and the Upper Mudstone Member (UMM), with the Lower Shale Member (LSM) of the Yinping Formation. The LPMM

and LCMM can be lithostratigraphically correlated to the PNMM and SRM, respectively.

Porous chert (Kametaka *et al.*, 2005; nectic chert in Zhu *et al.*, 1999 and Kametaka *et al.*, 2002) is a remarkable rock in the Gufeng Formation in the Lower Yangtze region. As its name suggests, the porous chert has abundant small cavities and shows a spongy structure compared to commonly-known normal chert characterized by being dense and non-porous. The bulk specific gravity of the porous chert is only 1.1–1.2 (Kametaka *et al.*, 2005). The porous chert is interbedded in the normal chert beds and is generally thicker than the normal chert beds.

Ma *et al.* (2016) and Shi *et al.* (2016) studied the lithostratigraphy of the Luojiaba and Macaojie sections near Jianshi, western Hubei, respectively. The Gufeng Formation in both sections consists primarily of siliceous rock with muddy rock. The lower parts of the Gufeng Formation in both sections are characterized by the dominance of dark-colored well-bedded chert (Figure 5D), whereas the upper parts are characterized by muddy rock including carbonates (Figure 5E). The Gufeng Formation in both sections is conformably covered by the limestone of the Wuxue Formation (Figure 5F). In these sections, the Gufeng Formation conformably overlies the Maokou Formation. Meanwhile, Chen *et al.* (2000) observed clastic rocks in the basal part of the Gufeng

Formation at Huangyan in Jianshi and concluded that the Gufeng Formation unconformably overlies the Maokou Formation there. The Bijiaoshan section contains thick carbonate rocks and are much thicker than the Luojiaba and Macaojie sections. This may indicate that the Bijiaoshan section represents shallower facies.

In short, the Gufeng Formation in the north margin is characterized by siliceous rocks with muddy rocks. Lithostratigraphy of the Gufeng Formation in the Lower Yangtze region comprises muddy rock, chert, and muddy rock with nodules in descending order whereas that in Middle Yangtze region consists of chert, muddy rock with carbonate, and muddy rock in ascending order.

South margin.—The bureaus of geology and mineral resources in Hunan, Jiangxi, Guangxi, and Guangdong established a hypostratotype section of the Gufeng Formation in each province: the Guanghuang section (27°17'E, 113°41'E) in You (Bureau of Geology and Mineral Resources of Hunan Province, 1997), the Majia'ao section (27°16'E, 113°53'E) in Lianhua (Bureau of Geology and Mineral Resources of Jiangxi Province, 1997), the Tongtianyan section in Liuzhou (Bureau of Geology and Mineral Resources of Guangxi Zhuang Autonomous Region, 1997), and the Youkeng section (24°32'E, 116°09'E) in Jianling (Bureau of Geology and Mineral Resources of

Guangdong Province, 1996).

These sections are characterized by much clastic rocks, such as carbonaceous mudstone, mudstone, and sandstone, with few cherts (Figure 4). The typical Gufeng Formation in the north margin, however, is characterized by siliceous rocks with muddy rocks as mentioned previously. We therefore believe that the “Gufeng Formation” in the south margin should be isolated from the Gufeng Formation and given another formation name.

Hu (2000) argued that the Gufeng Formation is also distributed over the south margin of the South China block based on discussions including fossil data. However, the Gufeng Formation is a lithostratigraphic unit. According to the International Stratigraphic Guide (Murphy and Salvador, 1999), “Lithostratigraphic units are defined and recognized by observable physical features and not by their inferred age, the time span they represent, inferred geologic history, or manner of formation”.

Radiolarian occurrences

The Gufeng Formation contains several macrofossils and microfossils, such as conodonts (e.g. Ching, 1960; Wang, 1995a; Kuwahara *et al.*, 2008a; Ma *et al.*, 2016),

foraminifers (e.g. Wang, 1993c), brachiopods (e.g. Jing and Hu, 1978; Xu *et al.*, 2004), sponge spicules (e.g. Kuwahara *et al.*, 2007b; Ito *et al.*, 2016e), mollusks (e.g. Xi, 1982; Kametaka *et al.*, 2009), and fish microremains (e.g. Kuwahara *et al.*, 2007b; Mao *et al.*, 2013).

Since the 1980s, radiolarian occurrences have been reported in the Gufeng Formation in several areas. On that basis, radiolarian zones have been constructed. Meanwhile, integrated radiolarian biostratigraphy has been compiled in Southwest Japan (Ishiga, 1986, 1990; Kuwahara *et al.*, 1998) and South China (Wang and Yang, 2007, 2011). The radiolarian zones of Southwest Japan, South China, the Gufeng Formation in the north margin, and the “Gufeng Formation” in the south margin are summarized in Figure 6.

Radiolarian biostratigraphy of the north margin

Bi (1981) found spherical radiolarians in thin sections from the Tianbaoshan section in southeast Nanjing, Jiangsu, which was probably the first report of radiolarians from the Gufeng Formation. Sheng and Wang (1985) derived radiolarian fossils from siliceous rocks in the Longtan area in Nanjing using a hydrofluoric acidic method and revealed radiolarian microphotographs via a scanning electron microscopy (SEM) for

the first time. Since these studies, several researchers have reported radiolarian occurrences in the Gufeng Formation in the Lower Yangtze region (Kong and Gong, 1986; Nagai and Zhu, 1992; Wang, 1993a, b, d, 1995b, c; Jiang *et al.*, 1994; Wang and Qi, 1995; Wang *et al.*, 1997; Nagai *et al.*, 1998; He *et al.*, 1999; Kuwahara *et al.*, 2007b; Kametaka *et al.*, 2009; Ito *et al.*, 2013a; Zhang *et al.*, in press). Sheng and Wang (1985) constructed the *Pseudoalbaillella scalprata*-*Ps. nanjingensis* and *Phanicosphaera mammilla* assemblage zones, in ascending order. Wang (1995c) found the first occurrence of *Follicucullus monacanthus* Ishiga and Imoto in the Gufeng Formation. Wang (1995b) discussed the age assignment of *F. monacanthus* and concluded that the *F. monacanthus* assemblage from the Gufeng Formation corresponds to that of the *F. monacanthus* zone of Japan and U.S.A. (Ishiga, 1986, 1990; Blome and Reed, 1992). Wang and Qi (1995) constructed the following assemblage zones: *Ps. fusiformis*-*Ps. longtanensis*, *F. monacanthus*, and *Ruzhencevispongus uralicus*-*F. scholasticus* in ascending order. Since the study by Wang and Qi (1995), the Gufeng Formation in the Lower Yangtze region has been divided into three radiolarian assemblage zones (e.g. He *et al.*, 1999; Kametaka *et al.*, 2009; Ito *et al.*, 2013a).

Feng (1992) discovered radiolarians in the Wuchan area, which was the first radiolarian report from the Gufeng Formation in the Middle Yangtze region. Since this

study, several researchers have reported radiolarian occurrences in this region (Feng and Zhong, 1994; Kuwahara *et al.*, 2007a, 2008a; Ma and Feng, 2012; Ma *et al.*, 2016; Shi *et al.*, 2016). Ma and Feng (2012) constructed the *Pseudoalbaillella globosa*, *Follicucullus monacanthus*, and *F. scholasticus* assemblage zones in ascending order.

The radiolarian zonation of the Lower Yangtze and Middle Yangtze regions are well correlated based on the lineage from *Pseudoalbaillella* Holdsworth and Jones to *Follicucullus* Ormiston and Babcock in the Guadalupian (e.g. Ishiga *et al.*, 1982; Wang and Yang, 2011; Zhang *et al.*, 2014; Ito *et al.*, 2015, 2016a).

In the Upper Yangtze region, fewer radiolarian studies have been performed compared to the Lower and Middle Yangtze regions. Kuwahara *et al.* (2007c) found *Copicyntra?* sp. and *Copiellintra?* sp. in the Gufeng Formation of the Mingyuexia 1 section in north Sichuan. They also found *Pseudoalbaillella* cf. *longtanensis* Sheng and Wang and *Ps. fusiformis* (Holdsworth and Jones) from the uppermost part of the Maokou Formation just below the Gufeng Formation in that section. Kuwahara *et al.* (2008b) found *Ps. fusiformis* and *Follicucullus* cf. *scholasticus* Ormiston and Babcock in the Chejiaba section in Chaotian, north Sichuan.

Radiolarian biostratigraphy of the south margin

The first occurrence of a radiolarian fossil from the “Gufeng Formation” in the south margin of the South China block was reported by Yao *et al.* (1993). They surveyed Late Paleozoic strata in Guizhou and Guangxi and obtained *Follicucullus scholasticus*, *Latentifistula* sp., and *Pseudoalbaillella* sp. from the “Gufeng Formation” in the Tongtianyan section in Guangxi. Wang and Li (1994) found *F. charveti* Caridroit and De Wever, *F. bipartitus* Caridroit and De Wever, and *Albaillella triangularis* Ishiga, Kito, and Imoto, corresponding to radiolarians of the *F. bipartitus*-*F. charveti* Assemblage Zone in Southwest Japan (Ishiga, 1986, 1990). Xian and Zhang (1998) researched the Bancheng area in Guangxi and constructed the following ascending radiolarian zones: *Ps. scalprata*, *A. sinuata*, *Ps. banchengensis*, *Ps. globosa*-*Ps. fusiformis*, *F. monacanthus*, and *F. scholasticus*. According to Xian and Zhang (1998), these zones correspond to radiolarian zones in Southwest Japan found by Ishiga (1986, 1990). However, judging from the text and figures of Xian and Zhang (1998), some of the radiolarian zones they constructed do not correspond directly to the radiolarian zones of Ishiga (1986, 1990). For example, *F. scholasticus*, which is a characteristic species of *F. scholasticus* Assemblage Zone of Ishiga (1986, 1990), occurred in the *Ps. globosa*-*Ps. fusiformis* and *F. monacanthus* zones of Xian and Zhang (1998). The *A. sinuata* Zone of Ishiga (1986, 1990) is a range zone defined by the occurrence of *A.*

sinuata. However, Xian and Zhang (1998) divided the *A. sinuata*-bearing strata of their section into the *A. sinuata* and *Ps. banchengensis* assemblage zones. Comparing the radiolarian zonation of Ishiga (1986, 1990) with that of Xian and Zhang (1998), the *Ps. longtanensis* Assemblage Zone, the *Ps. globosa* Assemblage Zone, and the *F. monacanthus* Range Zone of Ishiga (1986, 1990) are absent in the section of Xian and Zhang (1998). The Qinfang area, including the Bancheng area, is known as an orogenic belt in the late Paleozoic (e.g. He *et al.*, 2014; Ke *et al.*, 2018), and the orogeny might cause the absence of some biozones.

Kuwahara *et al.* (2004) found *Albaillella yamakitai* from the Tongtanyan section, Guangxi, which is a characteristic species of the *Follicucullus charveti*-*A. yamakitai* Assemblage Zone. It has been proposed that the base of the Lopingian of the radiolarian biostratigraphy is defined by the first occurrence of *A. yamakitai* or just below it (e.g. Mitsumura and Kamata, 2009; Nishikane *et al.*, 2011). This indicates that the top of the “Gufeng Formation” in Guangxi should reach the uppermost Guadalupian.

Wang and Yang (2003) found *Pseudoalbaillella fusiformis* and *Ps. globosa* in the Renhua area in Shaoguan, Guangdong. They constructed the *Ps. Fusiformis*-*Ps. globosa* Zone corresponding to the *Ps. Fusiformis*-*Ps. longtanensis* Zone in the Lower Yangtze region.

Stratigraphic change of radiolarian fauna

Stratigraphic change of radiolarian fauna has been clearly recognized in the Gufeng Formation in the north margin. Sheng and Wang (1985) divided radiolarians from the Longtan area into two faunas in ascending order: *Pseudoalbaillella* and *Phanicosphaera*. Kametaka *et al.* (2009) showed that *Albaillellaria* decrease but *Latentifistularia* increase upward in the Anmenkou section. Ito *et al.* (2013a) also recognized a similar pattern in the Liuhuang section. In the Middle Yangtze regions, *Albaillellaria* occurs in the lower parts of the Gufeng Formation, but it rarely occurs in the upper part (Shi *et al.*, 2016).

The radiolarian componential ratio at the order level (*Albaillellaria*, *Latentifistularia*, *Entactinaria*, and *Spumellaria*) had been used as an indicator of the water-depth (e.g. Kozur, 1993; Kuwahara, 1999). Kametaka *et al.* (2009) referred to these studies; however, they pointed out that the chief cause of the radiolarian faunal change in the Gufeng Formation in the Anmenkou section is not a water-depth change. This is because there are no remarkable lithological changes in the Anmenkou section. In other words, the water-depth of the Anmenkou section had to change by several hundreds of meters if water-depth changes are a critical factor for the faunal change.

Recently, Xiao *et al.* (2017) studied the latest Permian radiolarian vertical

distribution based on a correspondence analysis. Their result suggests that radiolarians exhibited different water depth at the species level even in the same order. Likewise, further studies on radiolarian vertical distributions in the middle Permian are necessary to understand the faunal change in the Gufeng Formation.

Remarkable radiolarian occurrences

In this section, we focus on remarkable radiolarian taxa and forms from primarily taxonomic and paleobiogeographic points of view, combined with our previous studies in other areas.

Longtanella Sheng and Wang.—*Longtanella* is likely an endemic genus of the Gufeng Formation (Figure 7A–C). *Longtanella* generally occurs in the cherts of the Gufeng Formation of the Lower Yangtze (Wang, 1993a; Kametaka *et al.*, 2009; Ito *et al.*, 2013a) and Middle Yangtze (Feng, 1992; Shi *et al.*, 2016) regions. This genus also occurs in the chert of the “Gufeng Formation” in the south margin (Xian and Zhang, 1998). Probable *Longtanella* occurs in the Sa Kao Formation in the Sra Kao suture zone in eastern Thailand (Saesaengseerung *et al.*, 2009) and the Kamiaso Unit in central Japan (Ito *et al.*, 2016d).

The taxonomy of *Longtanella*, however, is currently debatable. Since Sheng and Wang (1985) described the genus, subsequent studies mainly in China have used this taxonomic name (e.g. Kozur and Mostler, 1989; Feng, 1992; Wang, 1993a; Kametaka *et al.*, 2009; Ito *et al.*, 2013a; Shi *et al.*, 2016; Xiao *et al.*, 2018). Meanwhile, several previous studies have synonymized this genus with *Pseudoalbaillella* (e.g. De Wever *et al.*, 2001; Nestell and Nestell, 2010) or *Parafollicucullus* Holdsworth and Jones (Noble *et al.*, 2017). *Longtanella* was diagnosed as “Shell smooth, straight, bilaterally symmetrical turritiformis. The shell wall divided into the spire, the turri-body and the turri-bottom composed of ring-like swollen segments. Last segment constrictive, with 4 flaps vertically extending downward” (Sheng and Wang, 1985, p. 179). According to the diagnosis, description, and holotype (Figure 8) of *Longtanella*, the genus seems to differ from *Pseudoalbaillella* and *Parafollicucullus* in lacking winged pseudothorax and having four flaps. Further detailed observations are needed to solve this taxonomic problem.

Ruzhencevispongos uralicus Kozur.—*Ruzhencevispongos uralicus* is a characteristic species of the upper part of the Gufeng Formation (e.g. Wang, 1993a, b, 1995c; Wang and Qi, 1995; Kametaka *et al.*, 2009; Ito *et al.*, 2013a). This species possesses a

triangular platy spongy shell (Kozur, 1980) (Figures 7D, 7E). Wang (1993d) examined the morphology of *R. uralicus* with samples from Chaohu and Hushan. He divided this species into three morphotypes, namely *triadiatus*, *subtriadiatus*, and *uralicus*. Even though *Ruzhencevispongos* commonly occurs in South China including in the Gufeng Formation, only few such occurrences have been reported in Southwest Japan. (e.g. Ishiga, 1990).

Hegleria Nazarov and Ormiston.—Occurrences of this genus have been reported worldwide, and this genus is seemingly a cosmopolitan taxon. However, there are similar outer shaped radiolarians in the Permian. Ito *et al.* (2017c) stated these radiolarians as “*Hegleria*-shaped” radiolarians. The inner structure is important to identify “*Hegleria*-shaped” radiolarians. *Hegleria* has multiple internal rays that connect a central sphere to the outer shell (e.g. Noble and Jin, 2010) (Figure 7F). Several previous studies in the Gufeng Formation have shown radiolarians having this structure (e.g. Wang, 1993a, b, 1995c; Wang and Qi, 1995; Kametaka *et al.*, 2009; Ito *et al.*, 2013a), indicating that *Hegleria* surely occurred in the Gufeng Formation.

Meanwhile, “*Hegleria*-shaped” radiolarians having a concentric shell, probably *Paracopicyntra* spp., have also discovered in the Gufeng Formation (e.g. Wang, 1993a,

1995b; Kametaka *et al.*, 2009). Further, Kametaka *et al.* (2009) reported “*Hegleria*-shaped” radiolarians having a discoidal form as *Hegleria* sp. aff. *H. mammilla*, which might be *Guiuva sashidai* Ito and Feng.

Pseudotormentus De Wever and Caridroit.—*Pseudotormentus* commonly occurs in deposits from the Panthalassa (e.g. De Wever and Caridroit, 1984; Blome and Reed, 1992, 1995; Mitsumura and Kamata, 2009; Nakae, 2011; Ito *et al.*, 2013b; Ito and Matsuoka, 2015a, b, 2016, 2018) but rarely occurs in deposits from the Paleotethys. The Gufeng Formation, comprised of deposits from the Paleotethys, has rarely yielded *Pseudotormentus*. Ito *et al.* (2016b) examined this uneven distribution of *Pseudotormentus* and pointed out the possibility that it is a facies genus of a Panthalassan equatorial warm current.

Kametaka *et al.* (2009), however, reported the common occurrence of *Pseudotormentus* in the Gufeng Formation in the Anmenkou section. On the basis of the hypothesis of Ito *et al.* (2016b), the Gufeng Formation of the Anmenkou section might have been locally affected by the Panthalassan equatorial warm current. Alternatively, other factors such as lithology might control the uneven distribution of *Pseudotormentus*.

Corythoecidae Nazarov.—No radiolarian fossils of the family Corythoecidae, characterized by a bilaterally symmetrical shell with a lateral foramen, have been discovered in the Gufeng Formation. Even though the occurrence of this family is very rare worldwide (De Wever *et al.*, 2001), Ito *et al.* (2017a) speculated that most species of the Corythoecidae occurred in phosphate-rich facies on the basis of a compilation of the previous occurrences and implied that this family preferred nutrient-rich conditions.

In the Lower Yangtze region, abundant phosphate nodules are contained in the lowermost part of the Gufeng Formation (e.g. Huang *et al.*, 1994; Kametaka *et al.*, 2009; Ito *et al.*, 2013a). Although no radiolarians have been obtained in this part of the Gufeng Formation, it might have the potential to yield species of the Corythoecidae if the hypothesis of Ito *et al.* (2017a) is correct.

Swollen type.—Dimorphic pairs, comprised of normal and swollen types, are known in some taxa of the family Albaillellaria (e.g. Ishiga, 1991; Ito and Matsuoka, 2015b, 2017; Ito, 2017; Ito *et al.*, 2018). The swollen type is characterized by having a swollen apical portion. Ishiga (1991) suggested the possibility that the dimorphic pairs reflect alternating generations.

Previously, no occurrences of the swollen type have been reported from the Gufeng Formation; however, Xian and Zhang (1998) have reported the swollen type of *Albaillella sinuata* Ishiga and Watase and *Pseudoalbaillella globosa* Ishiga and Imoto in the “Gufeng Formation” in the Bancheng area.

Conjoined or malformed radiolarians.—Conjoined or malformed radiolarians are known in both living and fossil specimens (e.g. De Wever, 1985; Anderson and Gupta, 1998; Itaki and Bjørklund, 2007; Dumitrica, 2013; Afanasieva and Amon, 2016; Ito *et al.*, 2017b). Afanasieva and Amon (2016) classified the conjoined or malformed radiolarians as two types: conjoined radiolarians that two or more individuals are joined together were named as a multiplicative type; radiolarians having double or branching structure (e.g. spine, arm) were named as a supplemental type.

Certain radiolarian specimen of the multiplicative type has never discovered from the Gufeng Formation so far. Meanwhile, a specimen from the Gufeng Formation in the Anmenkou section shown by Kametaka *et al.* (2009, p. 117, fig. 7.18, identified as *Ishigaum* sp.) possesses branching arm. The specimen can be classified as the supplemental type of Afanasieva and Amon (2016).

Age assignments

The age of the Gufeng Formation has been calibrated primarily based on fossil data. Ching (1960) discovered conodont *Jinogondolella nankingensis* (Ching) in the basal mudstones of the Gufeng Formation in the Longtan area, Jiangsu. Subsequent studies have found this species in the basal mudstones of the Gufeng Formation in the Lower Yangtze region (e.g. Wang, 1995a; Kametaka *et al.*, 2009). *Jinogondolella nankingensis* is an index species of the Roadian (Ogg *et al.*, 2016). Meanwhile, Kametaka *et al.* (2009) also found ammonoids *Shouchangoceras* sp. and *Erinoceras* sp. from mudstones of the lower PNMM in the Anmenkou section, Anhui. The ranges of *Shouchangoceras* and *Erinoceras* are contained in the Wordian and the Roadian–Wordian, respectively (Zhao and Zheng, 1977); therefore, Kametaka *et al.* (2009) concluded that the age of the lower PNMM is Wordian and that of the SRM is Wordian or younger.

Ma *et al.* (2016) correlated conodont and radiolarian biozones in the Luojiaba section in west Hubei. They concluded that the *Pseudoalbaillella globosa*, *Follicucullus monacanthus*, and *F. scholasticus* radiolarian zones could be directly correlated with the *Jinogondolella nankingensis gracilis*, *J. aserrata*, and *J. postserrata* conodont zones.

In addition to the fossil data, Zhu *et al.* (2013) reported zircon U–Pb ages of

272.0±5.5 Ma (MSWD = 2.6) and 271.5±3.3 Ma (MSWD = 1.7) separated from volcanic ash beds within the basal shale strata of the Gufeng Formation in the Chaohu area, Anhui. The age of the boundary between the Kungurian (Cisuralian) and Roadian is 272.3 Ma according to the 2016 geologic time scale (Ogg *et al.*, 2016), so that the zircon U–Pb ages can be correlated to the lowermost part of the Roadian, Guadalupian. Zhang *et al.* (2010) correlated the conodont and radiolarian biozones across the Cisuralian-Guadalupian boundary and concluded that the base of Roadian coincides with the first occurrence of *Ps. globosa*. Consequently, the zircon U–Pb ages are consistent with the fossil data, even though the taxonomy of *Pseudoalbaillella globosa* is debatable (Ito *et al.*, 2016c).

In short, the Gufeng Formation is nearly consistent with the Guadalupian based on the conodonts and zircon U–Pb ages. In particular, the base of the Gufeng Formation should be near the boundary between the Cisuralian and Guadalupian. However, the top of the Gufeng Formation is uncertain because of fewer fossil occurrences and lack of numerical age dating.

Geochemical characteristics

Geochemical studies on the Gufeng Formation have increased since the 1990s (e.g. Lu and Qu, 1990; Xia *et al.*, 1995; Yang and Feng, 1997). Siliceous rocks within the Maokou Formation, which is a heterotopic facies of the Gufeng Formation, have also been geochemically investigated (e.g. Zhou *et al.*, 2009).

Origin of siliceous rock

Most previous studies have concluded that the siliceous rocks of the Gufeng Formation on the north margin are biogenic. The Al–Fe–Mn diagram defined by Adachi *et al.* (1986) and Yamamoto (1987) has generally been used in speculations of the origin of chert in the Paleozoic and Mesozoic (e.g. Du *et al.*, 2007; Ito *et al.*, 2016f). Most researchers have applied this diagram to the cherts of the Gufeng Formation. Most previous studies date the analyzed samples in the biogenesis field, or near it, in the Al–Fe–Mn diagram but not near the hydrothermal field (e.g. Kametaka *et al.*, 2005; Wang *et al.*, 2008; Yang and Yao, 2008; Han *et al.*, 2014). Meanwhile, a few studies have indicated the influence of hydrothermal activities in the Lower Yangtze region (Xia *et al.*, 1995; Xie *et al.*, 2013). Consequently, in general, a hydrothermal component

barely contributed little to the composition of the rocks in the Gufeng Formation except in a few areas.

In contrast to the Yangtze region, a scattering of studies on the “Gufeng Formation” in the south margin indicate that the cherts are likely hydrothermal. Fu *et al.* (2004) investigated the “Gufeng Formation” in Shaoyang in Hunan and showed that the cherts plot in the hydrothermal field on the Al–Fe–Mn diagram. Shi *et al.* (2016) pointed out the differences in the chert formation processes between the north and south margins. Further research is necessary to interpret the origin of siliceous rock of the “Gufeng Formation” on south margin because of fewer studies compared to the studies on the Gufeng Formation in the north margin.

Origin of organic matter

In contrast to the studies on the origin of siliceous rocks, a few studies have dealt with the origin of organic matter within the Gufeng Formation. Takebe *et al.* (2007) investigated organic matter in the black chert of the Anmenkou section using the van Krevelen diagram (atomic H/C ratio versus atomic O/C ratio). This diagram shows four types of kerogen according to the origin of the organic matter (van Krevelen, 1961). The analyzed data of Takebe *et al.* (2007) plotted in the field of type IV indicating reworked

and highly oxidized organic material. Oxidic weathering can cause an increase in the atomic O/C ratio of kerogen (e.g. Petsch *et al.*, 2000). However, due to several geochemical results, such as the small variation of the atomic O/C ratio, S enrichment, and the inclusion of pyrites, Takabe *et al.* (2007) concluded that the black chert was not affected by oxidic weathering and originated from terrestrial or reworked organic matter.

However, considering the high degree of maturity estimated in a subsequent study, the analyzed data by Takebe *et al.* (2007) were likely not primary. In general, kerogens subjected to increasing thermal stress move toward the lower left corner of the van Krevelen diagram (Hunt, 1995). This tendency has been reported by laboratory experiments (e.g. Kotarba and Lewan, 2004; Lewan and Kotarba, 2014; Takahashi and Suzuki, 2017). According to the geochemical works by Du *et al.* (2015), the vitrinite reflectance (VR %) and Rock-Eval T_{max} of siliceous and muddy rocks in the Gufeng Formation have a range of 1.2–2.0% and 460–560°C, respectively. These values indicate a high degree of maturity. Under such a high degree of maturity, the van Krevelen diagram is not applicable to estimate of the origin of the organic matter.

The supply of terrestrial material to the Gufeng Formation is irrefutable due to several observed facts as mentioned in the previous chapter, such as the presence of intercalated clastics within the Gufeng Formation. The origin of the organic matter in

the Gufeng Formation is debatable; therefore, further data such as the stable carbon isotopic composition of the organic matter are needed to discuss its origin.

Sedimentary environment

There had been speculations as to the sedimentary environment of the Gufeng Formation based on the lithology and radiolarian faunal comparison. Kong and Gong (1987) postulated that the siliceous rocks of the Gufeng Formation were formed in a deep-sea environment resulting from rift faulting because the radiolarian fauna is similar to the Japanese assemblage. Zhu (1989) concluded that the Gufeng Formation was deposited in an intraplatformal environment. Since the 1990s, a geochemical approach to speculations of the sedimentary environment of the Gufeng Formation has been studied. Such studies have generally concluded that the Gufeng Formation formed in a continental shelf environment (e.g. Kametaka *et al.*, 2005).

Murray (1994) placed the pelagic and continental margin fields in the $\text{Fe}_2\text{O}_3/\text{TiO}_2$ versus $\text{Al}_2\text{O}_3/(\text{Al}_2\text{O}_3+\text{Fe}_2\text{O}_3)$ diagram on the basis of chert from continental margin, pelagic, and ridge-proximal environments, ranging from the Early Paleozoic to the Neogene. To speculate about the sedimentary environment, the diagram has been used (e.g. Hara *et al.*, 2010; Ito *et al.*, 2016f; Kemkin *et al.*, 2017). Most of the analyzed

samples from the Gufeng Formation plot in the continental margin field in the diagram (e.g. Kametaka *et al.*, 2005; Wang *et al.*, 2008; Yang and Yao, 2008; Han *et al.*, 2014).

Redox conditions and changes

Redox conditions and changes during the deposition of the Gufeng Formation have been studied in detail, especially in the Lower Yangtze (Kametaka *et al.*, 2005) and the Middle Yangtze (Shi *et al.*, 2016) regions. In short, the redox condition of the Gufeng Formation in the Lower Yangtze region changed from aerobic to suboxic to anoxic, whereas that in the Middle Yangtze region changed from oxic to anoxic to suboxic. In this chapter, we focus on the Lower and Middle Yangtze regions based on the discussion and speculation of the previous studies.

The Anmenkou section in the Lower Yangtze region is divided into the PNMM and SRM in ascending order (Kametaka *et al.*, 2005). The lowermost mudstone of the PNMM contains glauconite pellets. Glauconites are generally formed under aerobic shallow water (e.g. Tucker, 1991; Prothero and Schwab, 1996). Consequently, the lower PNMM was formed under such conditions.

The upper PNMM contains abundant phosphate nodules. Phosphate nodules are generally formed under oxygen-minimum zones during low clastic supply on the outer

shelf and upper slope (e.g. Trucker, 1991; Prothero and Schwab, 1996; Trappe, 1998).

Kametaka *et al.* (2005) therefore concluded that the upper PNMM is formed under suboxic conditions.

Kametaka *et al.* (2005) concluded that the SRM of the Anmenkou sections were formed in a continental margin under anoxic conditions on the basis of several features of the chert, such as high Mo, Ni, Cu, and V and extremely low Mn contents. These elements are used as an indicator of the redox condition of the sediments. For example, Mo is scavenged by Mn-oxides under oxic conditions (Shimmield and Price, 1986); Mo is dissolved and remobilized under suboxic conditions (Chaillou *et al.*, 2002; McManus *et al.*, 2002; Shimmield and Price, 1986); and Mo forms strong associations with sulfide, concentrating in deposits (Vorlicek *et al.*, 2004). Nickel and Cu also concentrate in sediments under reducing conditions (Calvert and Pedersen, 1993). Vanadium in deposits primarily originates from continental flux or inclusion in seawater. The V and U originating from continental flux decrease in proportion to the Al concentration. Conversely, the seawater-derived V in sediments depends on the redox condition of the deposits (Hori *et al.*, 2000). Manganese generally deposits as Mn-oxide (MnO_2) under oxidizing conditions, whereas it becomes Mn^{2+} and readily secedes from deposits under reducing conditions (Calvert and Pedersen, 1993).

In the Middle Yangtze region, Shi *et al.* (2016) subdivided the Maocaojie section into three units. The lower unit consists primarily of siliceous rocks; the middle unit is composed of limestone, carbonous mudstone, and muddy chert; and the upper part consists of muddy chert, siliceous mudstone, and calcareous mudstone. The geochemical results by Shi *et al.* (2016) indicated that the lower, middle, and upper units were formed under oxic, suboxic, and anoxic conditions, respectively. On the basis of comparison to the stratigraphic change of the radiolarian component, Shi *et al.* (2016) concluded that the widespread anoxia was promoted by the elevated primary productivity.

Concluding remarks

In this article, we reviewed the lithostratigraphy, radiolarian occurrences with biostratigraphy, and geochemical characteristics of the Gufeng Formation. Here we summarize the lithostratigraphy and redox conditions of the Gufeng Formation of the Anmenkou and Maocaojie sections as representative examples of the Lower and Middle Yangtze regions, respectively (Figure 9).

In the Anmenkou section of the Lower Yangtze region, the dominant rock facies of

the Gufeng Formation are muddy rock with nodules, chert, and muddy rock in ascending order. The former is called the PNMM; the latter two are called the SRM by Kametaka *et al.* (2002, 2005). Based on the lithology, the lower PNMM formed under aerobic shallow water and the upper PNMM formed under suboxic conditions (Kametaka *et al.*, 2005). The geochemical characteristics of the SRM indicate that it was formed under anoxic conditions (Kametaka *et al.*, 2005).

In the Maocaojie section of the Middle Yangtze region, the dominant rock facies of the Gufeng Formation are chert, muddy rock with carbonate, and muddy rock (Shi *et al.*, 2016). The geochemical results indicate that these parts were formed under oxic, anoxic, and suboxic conditions, respectively (Shi *et al.*, 2016).

In contrast to the Lower and Middle Yangtze regions, fewer studies have focused on the Gufeng Formation in the Upper Yangtze region (e.g. Kuwahara *et al.*, 2007c, 2008a; Zhou *et al.*, 2009). In the recent decade, upper Capitanian carbonates of the Maokou Formation in the Chaotian section in northern Sichuan have been studied in detail (e.g. Lai *et al.*, 2008; Saitoh *et al.*, 2013, 2014; Chen *et al.*, 2011). Further studies on stratigraphic change in the lithostratigraphy, radiolarian biostratigraphy, and geochemical characteristics of the Gufeng Formation and chert beds within the carbonates of the Maokou Formation in the Upper Yangtze region should reveal the

paleoenvironmental development on the north margin of the South China block.

Acknowledgements

We thank Weihong He (China University of Geosciences, Wuhan) and Masao Kametaka (Dia Consultant Co. Ltd.) for critical review on this manuscript and valuable suggestions. Comments given by Associate Editor, Toshiyuki Kurihara (Niigata University), improved greatly this manuscript. We are most grateful to the following students and PhD students of China University of Geosciences, Wuhan, for kind help for fieldwork of the Gufeng Formation in Chaohu (Anhui) and Jianshi (Hubei) and collecting and reading the Chinese articles: Lei Zhang, Yong Lei, Qing Hu, Lei Shi, Jun Shen, Wenchao Cao, Qiangfen Ma, Qi Wu, Hailong Gao, and others. This study has been financially supported by NSFC (40839903 and 40921062) and “111” Project (B08030) of China University of Geosciences, Wuhan and “Niigata University’s Scholarship Program for Graduate School Students conducting Research Abroad”.

References

- Adachi, M., Yamamoto, K. and Sugisaki, R., 1986: Hydrothermal chert and associated siliceous rocks from the northern Pacific: their geological significance as indication of ocean ridge activity. *Sedimentary Geology*, vol. 47, p. 125–148.
- Afanasieva, M. S. and Amon, E. O., 2016: Deviations in skeletons of radiolarians. *Paleontological Journal*, vol. 50, p. 1529–1543.
- Anderson, O. R. and Gupta, S. M., 1998: Evidence of binary division in mature central capsules of a collosphaerid colonial radiolarian, implications for shell ontogenetic patterns in modern and fossil species. *Palaeontologia Electronica*, vol. 1, p. 1–13.
- Beauchamp, B. and Baud, A., 2002: Growth and demise of Permian biogenic chert along northwest Pangea: evidence for end-Permian collapse of thermohaline circulation. *Palaeogeography, Palaeoclimatology, Palaeoecology*, vol. 184, p. 37–63.
- Bi, Z. Q., 1981: Discovery of radiolarian fossils from the Permian Gufeng Formation and Dalong Formation near Nanjing area and its significance. *Nanjing Institute of Geology and Mineral Resources, Academia Sinica*, vol. 3, p. 111–112. (in Chinese)
- Blome, C. D. and Reed, K. M., 1992: Permian and Early (?) Triassic radiolarian faunas

from the Grindstone terrane, central Oregon. *Journal of Paleontology*, vol. 66, p. 351–383.

Blome, C. D. and Reed, K. M., 1995: Radiolarian biostratigraphy of the Quinn River Formation, Black Rock terrane, north–central Nevada: Correlations with eastern Klamath terrane geology. *Micropaleontology*, vol. 41, p. 49–68.

Bureau of Geology and Mineral Resources of Anhui Province, 1997: *Stratigraphy (Lithostratic) of Anhui Province*, 296 p. CUG Press, Wuhan. (in Chinese)

Bureau of Geology and Mineral Resources of Guangdong Province, 1996: *Stratigraphy (Lithostratic) of Guangdong Province*, 264 p. CUG Press, Wuhan. (in Chinese)

Bureau of Geology and Mineral Resources of Guangxi Zhuang Autonomous Region, 1997: *Stratigraphy (Lithostratic) of Guangxi Zhuang Autonomous Region*, 310 p. CUG Press, Wuhan. (in Chinese)

Bureau of Geology and Mineral Resources of Hubei Province, 1996: *Stratigraphy (Lithostratic) of Hubei Province*, 284 p. CUG Press, Wuhan. (in Chinese)

Bureau of Geology and Mineral Resources of Hunan Province, 1997: *Stratigraphy (Lithostratic) of Hunan Province*, 292 p. CUG Press, Wuhan. (in Chinese)

Bureau of Geology and Mineral Resources of Jiangsu Province, 1997: *Stratigraphy (Lithostratic) of Jiangsu Province*, 288 p. CUG Press, Wuhan. (in Chinese)

- Bureau of Geology and Mineral Resources of Jiangxi Province, 1997: *Stratigraphy (Lithostratic) of Jiangxi Province*, 381 p. CUG Press, Wuhan. (in Chinese)
- Bureau of Geology and Mineral Resources of Sichuan Province, 1997: *Stratigraphy (Lithostratic) of Sichuan Province*, 417 p. CUG Press, Wuhan. (in Chinese)
- Bureau of Geology and Mineral Resources of Zhejiang Province, 1996: *Stratigraphy (Lithostratic) of Zhejiang Province*, 236 p. CUG Press, Wuhan. (in Chinese)
- Calvert, S. E. and Pedersen, T. F., 1993: Geochemistry of recent oxic and anoxic marine sediments: implications for the geological record. *Marine Geology*, vol. 113, p. 67–88.
- Chaillou, G., Anschutz, P., Lavaux, G., Schafer, J. and Blang, G., 2002: The distribution of Mo, U, and Cd in relation to major redox species in muddy sediments of the Bay of Biscay. *Marine Chemistry*, vol. 80, p. 41–59.
- Chen, B., Joachimski, M. M., Sun, Y. D., Shen, S. Z. and Lai, X. L., 2011: Carbon and conodont apatite oxygen isotope records of Guadalupian-Lopingian boundary sections: climatic or sea-level signal? *Palaeogeography, Palaeoclimatology, Palaeoecology*, vol. 311, p. 145–153.
- Chen, L. D., Li, Z. H., Huang, Z. X., Zhang, K. M. and Duan, W. J., 2000: Discovery of the clastic rocks in the basal part of the Permian Gufeng Formation in Huangyan,

- Jianshi, Hubei and its significance. *Journal of Stratigraphy*, vol. 24, p. 207–209. (in Chinese with English abstract)
- Ching, Y., 1960: Conodonts from the Kufeng Suite (Formation) of Lungtan, Nanking. *Acta Paleontologica Sinica*, vol. 8, p. 242–248. (in Chinese with English abstract)
- De Wever, P., 1985: Sur l'existence, dès le Paléozoïque, de radiolaires siamois. *Revue de Paléobiologie*, vol. 4, p. 111–116. (in French with English abstract)
- De Wever, P. and Caridroit, M., 1984: Description de Quelques Nouveaux Latentifistulidea (Radiolaires Polycystines) Paleozorques du Japon. *Revue de Micropaléontologie*, vol. 27, p. 98–106. (in French with English abstract)
- De Wever, P., Dumitrica, P., Caulet, J. P., Nigrini, C. and Caridroit, M., 2001: *Radiolarians in the Sedimentary Record*, 533 p. Gordon and Breach Science Publishers, Singapore.
- Du, Y. S., Zhu, J., Gu, S. Z., XU, Y. J. and Yang, J. H., 2007: Sedimentary geochemistry of the Cambrian-Ordovician cherts: Implication on archipelagic ocean of North Qilian orogenic belt. *Science in China Series D, Earth Sciences*, vol. 50, p. 1628–1644.
- Du, X. B., Song, X. D., Zhang, M. Q., Lu, Y. C., Lu, Y. B., Chen, P., Liu, Z. H. and Yang, S. A., 2015: Shale gas potential of the Lower Permian Gufeng Formation in

- the western area of the Lower Yangtze Platform, China. *Marine and Petroleum Geology*, vol. 67, p. 526–543.
- Dumitrica, P., 2013: Siamese twins and twin-like skeletons in Mesozoic Polycystine Radiolaria. *Revue de Micropaléontologie*, vol. 56, p. 51–61.
- Feng, Q. L., 1992: Permian and Triassic radiolarian biostratigraphy in south and southwest China. *Journal of China University of Geosciences*, vol. 3, p. 51–62.
- Feng, Q. L. and Zhong, C. T., 1994: A new research method of sedimentary environments of bedded siliceous rocks, with discussing on sedimentary environments of the Gufeng Formation in Wuchan area, Hubei. *Sedimentary Facies and Palaeogeography*, vol. 14, p. 45–55. (in Chinese with English abstract)
- Fu, W., Zhou, Y. Z., Yang, Z. J., He, J. G., Zhang, C. B. and Yang, H. S., 2004: Petrogenesis of the bedded chert from the Gufeng Formation and its implications to Early Permian geodynamic background in South China. *Bulletin of Mineralogy, Petrology and Geochemistry*, vol. 23, p. 292–300. (in Chinese with English abstract)
- Golonka, J. and Ford, D., 2000: Pangean (Late Carboniferous–Middle Jurassic) paleoenvironment and lithofacies. *Palaeogeography, Palaeoclimatology, Palaeoecology*, vol. 161, p. 1–34.

- Han, Z. Z., Xiao, N., Li, A. L., Gao, F., Xu, C. L. and He, Y. Y., 2014: Geochemistry and sedimentary environments analysis of siliceous rocks from the Gufeng Formation of Lower Permian in Chaohu Region, Anhui Province. *Periodical of Ocean University of China*, vol. 44, p. 78–99. (in Chinese with English abstract)
- Hara, H., Kurihara, T., Kuroda, J., Adachi, Y., Kurita, H., Wakita, K., Hisada, K., Charusiri, P., Charoentitirat, T. and Chaodumrong, P., 2010: Geological and geochemical aspects of a Devonian siliceous succession in northern Thailand: Implications for the opening of the Paleo-Tethys. *Palaeogeography, Palaeoclimatology, Palaeoecology*, vol. 297, p. 452–464.
- He, W. H., Wu, S. B., Zhang, K. X. and Bu, J. J., 1999: Classification of radiolarian fossil zones and environmental analysis of Gufeng Formation in Lower Yangtze region. *Jiangsu Geology*, vol. 23, p. 17–23. (in Chinese with English abstract)
- He, W. H., Tang, T. T., Yue, M. L., Deng, J. F., Pan, G. T., Xing, G. F., Luo, M. S., Xu, Ya. D., Wei, Y., Zhang, Z. Y., Xiao, Y. F. and Zhang, K. X., 2014: Sedimentary and tectonic evolution of Nanhuan-Permian in South China. *Earth Science—Journal of China University of Geoscience*, vol. 39, p. 929–953. (in Chinese with English abstract)
- Hein, J. R., 2004: The Permian Earth. In, Hein, J. R., ed., *Life Cycle of the Phosphoria*

- Formation: From Deposition to Post-Mining Environment*, p. 3–17. Handbook of Exploration and Environment Geochemistry, Vol. 8. Elsevier, Amsterdam.
- Hori, S. R., Higuchi, Y. and Fujiki, T., 2000: Chemical compositions and their environmental records of bedded cherts from accretionary complex. *Memoirs of Geological Society of Japan*, no. 55, 43–59 (in Japanese with English abstract).
- Hu, S. Z., 2000: New consideration of Gufeng Formation by stratigraphy check up. *Volcanology & Mineral Resources*, vol. 21, p. 63–68. (in Chinese with English abstract)
- Huang, M. K., Qing, C. S. and Xing, Y. C., 1994: Petrological characteristics and depositional environment analysis of Permian system in Zuoshan, Susong County, Anhui province. *Journal of China University of Mining and Technology*, vol. 23, p. 89–94. (in Chinese with English abstract)
- Hunt, J. M., 1995: The source rock. *Petroleum Geochemistry and Geology*, Second Edition, p. 322–394. W. H. Freeman and Company, New York.
- Isbell, J. L., Henry, L. C., Gulbranson, E. L., Limariano, C. O., Fraiser, M. L., Koch, Z. J., Ciccio, P. L. and Dineen, A. A., 2011: Glacial paradoxes during the late Paleozoic ice age: Evaluating the equilibrium line altitude as a control on glaciation. *Gondwana Research*, vol. 22, p. 1–19.

- Ishiga, H., 1986: Late Carboniferous and Permian radiolarian biostratigraphy of southwest Japan. *Journal of Osaka City University*, vol. 29, p. 89–100.
- Ishiga, H., 1990: Paleozoic radiolarians. In, Ichikawa, K., Mizutani, S., Hara, I., Hada, S. and Yao, A. eds., *Pre-Cretaceous Terrane of Japan*, p. 285–295. Publication of IGCP project No. 224: Pre-Jurassic Evolution of Eastern Asia, Osaka.
- Ishiga, H., 1991: “Dimorphic pairs” of Albaillellaria (Late Paleozoic radiolaria), Japan. *Memoir of the Faculty of Science, Shimane University*, vol. 25, p. 119–129.
- Ishiga, H., Kito, T. and Imoto, N., 1982: Middle Permian radiolarian assemblages in the Tamba district and an adjacent area, southwest Japan. *Earth Science (Chikyu Kagaku)*, vol. 36, p. 272–281.
- Isozaki, Y., Kawahata, H. and Minoshima, K., 2007a: The Capitanian (Permian) Kamura cooling event: The beginning of the Paleozoic–Mesozoic transition. *Palaeoworld*, vol. 16, p. 16–30.
- Isozaki, Y., Kawahata, H. and Ota, A., 2007b: A unique carbon isotope record across the Guadalupian–Lopingian (Middle–Upper Permian) boundary in mid-oceanic paleo-atoll carbonates: The high-productivity “Kamura event” and its collapse in Panthalassa. *Global and Planetary Change*, vol. 55, p. 21–38.
- Itaki, T. and Bjørklund, K. R. 2007: Conjoined radiolarian skeletons (Actinommidae)

- from the Japan Sea sediments. *Micropaleontology*, vol. 53, p. 371–389.
- Ito, T., 2017: Dimorphism of Permian Albaillellaria: Reproductive strategy and its development of fossil radiolaria. *Radiolaria*, vol. 40, p. 60–61.
- Ito, T., Feng, Q. L. and Matsuoka, A., 2013a: Radiolarian faunal change in the Middle Permian Gufeng Formation in the Liuhuang section, Chaohu, South China. *Science Reports of Niigata University (Geology)*, no. 28, p. 39–49.
- Ito, T., Feng, Q. L. and Matsuoka, A., 2015. Taxonomic significance of short forms of middle Permian *Pseudoalbaillella* Holdsworth and Jones, 1980 (Follicucullidae: Radiolaria). *Revue de Micropaléontologie*, vol. 58, p. 3–12.
- Ito, T., Feng, Q. L. and Matsuoka, A., 2016a. Possible boundaries between *Pseudoalbaillella* and *Follicucullus* (Follicucullidae, Albaillellaria, Radiolaria): An example of morphological information from fossils and its use in taxonomy. *Forma*, vol. 31, p. 7–10.
- Ito, T., Feng, Q. L. and Matsuoka, A., 2016b: Uneven distribution of *Pseudotormetus* De Wever et Caridroit (Radiolaria, Protozoa): Provincialism of a Permian planktonic microorganism. *Acta Geologica Sinica (English Edition)*, vol. 90, p. 1598–1610.
- Ito, T., Feng, Q. L. and Matsuoka, A., 2016c: A new subspecies and morphological variation of *Pseudoalbaillella globosa* (Middle Permian radiolaria). *Acta*

Micropalaontologica Sinica, vol. 33, p. 436–443.

Ito, T., Gu, S. Z., Ai, Y. and Feng, Q. L., 2017a: A new genus of the Corythoecidae (Paleozoic Radiolaria) from the Changhsingian (uppermost Permian) Dalong Formation in southern Guizhou, South China. *Paleontological Research*, vol. 21, p. 131–137.

Ito, T., Kitagawa, Y., and Matsuoka, A., 2016d: Middle and Late Permian radiolarians from chert blocks within conglomerates of the Kamiaso Unit of the Mino terrane in Gifu Prefecture, central Japan. *Journal of the Geological Society of Japan*, vol. 122, p. 249–259. (in Japanese with English abstract)

Ito, T., Kitagawa, Y. and Matsuoka, A., 2017b: An aberrant bi-apical *Follicucullus* (Albaillellaria) from the late Guadalupian (Middle Permian), with the possible oldest evidence on double malformation in radiolarians. *Journal of Micropalaeontology*, vol. 36, p. 222–223

Ito, T. and Matsuoka, A., 2015a: Imbricate structure of the Permian Yoshii Group in the Otakeyama area, Okayama Prefecture, southwest Japan. *Frontiers of Earth Science*, vol. 9, p. 152–163.

Ito, T. and Matsuoka, A., 2015b: Swollen type of *Albaillella sinuata* Ishiga and Watase (Permian radiolaria) from a chert boulder in Ie-jima Island, Okinawa Prefecture,

- Japan. *News of Osaka Micropaleontologists (NOM), Special Volume*, no. 15, p. 207–217.
- Ito, T. and Matsuoka, A., 2016: Ductilely deformed cherts within intrusive sandstones of the Yoshii Group of the Akiyoshi terrane in Southwest Japan: Consolidation time of Permian cherts. *News of Osaka Micropaleontologists (NOM), Special Volume*, no. 16, p. 95–104.
- Ito, T. and Matsuoka, A., 2017: Permian–Cretaceous radiolarians from Ie Island, Okinawa Prefecture, Japan. *Science Reports of Niigata University (Geology)*, no. 32 (supplement), p. 125–136.
- Ito, T. and Matsuoka, A., 2018: Lithology and radiolarian age of the Ryokami-yama Chert Formation in eastern Mt. Ryokami: Possible décollement zone in Permian pelagic sequence in mid-Mesozoic accretionary complexes of Southwest Japan. *Island Arc*, vol. 27, e12273.
- Ito, T., Matsuoka, A. and Feng, Q. L., 2016e: Siliceous sponge spicules from the Permian Gufeng Formation in Wuhan, Hubei Province, China. *News of Osaka Micropaleontologists (NOM), Special Volume*, no 16, p. 105–111.
- Ito, T., Matsuoka, A. and Feng, Q. L., 2018: Morphological characteristics of a dimorphic pair of *Albaillella sinuata* Ishiga and Watase: Dimorphism in the lineage

- of the Permian Albaillellaria (Radiolaria). *Island Arc*, vol. 27, e12271.
- Ito, T., Qian, X. and Feng, Q. L., 2016f: Geochemistry of Triassic siliceous rocks of the Muyinhe Formation in the Changning-Menglian belt of Southwest China. *Journal of Earth Science*, vol. 27, p. 403–411.
- Ito, T., Zhang, L., Feng, Q. L. and Matsuoka, A., 2013b: Guadalupian (Middle Permian) radiolarian and sponge spicule faunas from the Bancheng Formation of the Qinzhou allochthon, South China. *Journal of Earth Science*, vol. 24, p. 145–156.
- Ito, T., Zhang, L., Zhang, M. H., Feng, Q. L. and Matsuoka, A., 2017c: *Guiuva sashidai* n. gen. n. sp., a probable colonial radiolaria from the Lopingian (Upper Permian) in South China. *Palaeoworld*, vol. 26, p. 631–637.
- Jiang, N. Y., Jia, R. F., Wang, Z. Y., Qi, D. L., Yu, Z. Y., He, Y. M., Chen, X. M., Zhao, L., Zhuo, E. J., Zhang, X. H., Zhu, Z. L. and Zhu, Z. Q., 1994: *Permian Palaeogeography and Geochemical Environment in Lower Yangtze Region, China*, 214 p. Petroleum Industry Press, Beijing.
- Jin, Y. G., Zhang, J. and Shang, Q. H., 1994: Two phases of the end-Permian mass extinction. In, Embry, A. F., Beauchamp, B. and Glass, D. J. eds., *Pangea: Global Environments and Resources*, p. 813–822. Memoir Canadian Society of Petroleum Geologists, no. 17.

- Jing, Y. G. and Hu, S. Z., 1978: Brachiopods of the Kuhfeng Formation in south Anhui and Nanking hills. *Acta Palaeontologica Sinica*, vol. 17, p. 101–127 (*in Chinese*).
- Kametaka, M., Nagai, H., Zhu, S. and Takebe, M., 2009: Middle Permian radiolarians from Anmenkou, Chaohu, Northeastern Yangtze platform, China. *Island Arc*, vol. 18, p. 108–125.
- Kametaka, M., Takebe, M., Nagai, H. and Zhu, S., 2002: Sedimentary petrography of the Middle Permian Gufeng Formation in the Chaohu area, Anhui Province, China. *Bulletin of Nagoya University Museum*, vol. 18, p. 57–66. (*in Japanese with English abstract*)
- Kametaka, M., Takebe, M., Nagai, H., Zhu, S. and Takayanagi, Y., 2005: Sedimentary environments of the Middle Permian phosphorite-chert complex from the northeastern Yangtze platform, China; the Gufeng Formation: a continental shelf radiolarian chert. *Sedimentary Geology*, vol. 174, p. 197–222.
- Ke, X., Zhang, Z. Y., Yang, J. H., Yao, H. Z., Wang, S. D. and He, W. H., 2018: Radiolarian and detrital zircon in the Upper Carboniferous to Permian Bancheng Formation, Qinfang Basin, and the geological significance. *Journal of Earth Science*, vol. 29, p. 594–606.
- Kemkin, I. V., Khanchuk, A. I. and Kemkina, R. A., 2017: Geochemical evidence for

- consecutive accretion of oceanic fragments: The example of the Samarka terrane, Sikhote-Alin. *Doklady Earth Sciences*, vol. 474, p. 524–529.
- Kobayashi, F., 1997: Middle Permian biogeography based on fusulinacean faunas. In, Ross, C. A., Ross, J. R. P. and Brenckle, P. L. eds., *Late Paleozoic Foraminifera; Their Biostratigraphy, Evolution, and Paleoecology; and the Mid-Carboniferous Boundary*, p. 73–76. Cushman Foundation for Foraminiferal Research, Special publication, no. 36.
- Kong, Q. Y. and Gong, Y. J., 1986: Discovery of radiolarian siliceous rocks in the Maokou Stage of the Lower Permian in Chaoxian, Anhui, and their geological significance. *Geological Review*, vol. 32, p. 505–511. (in Chinese with English abstract)
- Kong, Q. Y. and Gong, Y. J., 1987: The formation environment of the Lower Permian radiolarian siliciliths in Jiangsu-Anhui region. *Oil and Gas Geology*, vol. 8, p. 86–89. (in Chinese with English abstract)
- Kotarba, M. J. and Lewan, M. D., 2004: Characterizing thermogenic coalbed gas from Polish coals of different ranks by hydrous pyrolysis. *Organic Geochemistry*, vol. 35, p. 615–646.
- Kozur, H., 1980: Ruzhencevispongidae, eine neue Spumellaria Familie aus dem oberen

- Kungurian (Leonardian) und Sakmarian des Vorurals. *Geologisch Paläontologische Mitteilungen Innsbruck*, Band 10, p. 235–242. (in German with English abstract)
- Kozur, H., 1993: Upper Permian radiolarians from the Sosio Valley area, western Sicily (Italy) and from the Uppermost Lamar Limestone of west Texas. *Jahrbuch der Geologischen Bundesanstalt*, Band 136, p. 99–123.
- Kozur, H. and Mostler, H., 1989: Radiolarien und Schwammskleren aus dem Unterperm des Vorurals. *Geologisch Paläontologische Mitteilungen Innsbruck, Sonderband*, Band 2, p. 146–275. (in German with English abstract)
- Kuwahara, K., 1999: Middle–Late Permian radiolarian assemblages from China and Japan. In, Yao, A., Ezaki, Y., Hao, W. and Wang, X. eds., *Biotic and Geological Development of the Paleo-Tethys in China*, p. 43–54. Peking University Press. Beijing.
- Kuwahara, K., Yao, A. and Yamakita, S., 1998: Reexamination of Upper Permian radiolarian biostratigraphy. *Earth Science (Chikyu Kagaku)*, vol. 52, p. 391–404.
- Kuwahara, K., Yao, A., Yao, J. X., Feng, S., Ji, Z. and Yao, H., 2007a: Middle Permian radiolarian biostratigraphy on the Gufeng Formation in the Songzi–Wufeng area, Hubei Province, China. *Journal of Geosciences, Osaka City University*, vol. 50, p. 55–66.

Kuwahara, K., Yao, A., Yao, J. X. and Wang, X., 2007b: Permian radiolarians from the Gufeng Formation of the Tongling area, Anhui Province, China. *Journal of Geosciences, Osaka City University*, vol. 50, p. 35–54.

Kuwahara, K., Yao, A., Yao, J. X. and Wu, Z. J., 2007c: Permian and Triassic radiolarian assemblages from the Yangzi platform (Part 10) –Radiolarian biostratigraphy of the Permian Gufeng Formation in the northern Sichuan, China. *Abstracts with Programs of the 2007 Annual Meeting of the Palaeontological Society of Japan*, p. 50. (in Japanese)

Kuwahara, K., Yao, A., Yao, J. X., Feng, S. N., Ji, Z. S. and Yao, H. Z., 2008a: Findings of Middle Permian radiolarian and conodont fossils from the Gufeng Formation of the Zigui area, Hubei Province, China. *Journal of Geosciences, Osaka City University*, vol. 51, p. 9–19.

Kuwahara, K., Yao, A., Yao, J. X. and Wu, Z. J., 2008b: Permian and Triassic radiolarian assemblages from the Yangzi platform (Part 11) –Radiolarian fossils from the Middle and Upper Permian in the northern Sichuan, China. *Abstracts with Programs of the 157th Regular Meeting of the Palaeontological Society of Japan*, p. 61. (in Japanese)

Kuwahara, K., Yao, A., Yao, J. X. and Li, J., 2004: Late Permian radiolarians and

- sponge spicules from the Tongtianyan section, Liuzhou, Guangxi, China. *Journal of Geosciences, Osaka City University*, vol. 47, p. 85–99.
- Lai, X. L., Wang, W., Wignall, P. B., Bond, D. P. G., Jiang, H. S., Ali, J. R., John, E. H., Sun, Y. D., 2008: Palaeoenvironmental change during the end-Guadalupian (Permian) mass extinction in Sichuan, China. *Palaeogeography, Palaeoclimatology, Palaeoecology*, vol. 269, p. 78–93.
- Lewan, M. D. and Kotarba, M. J., 2014: Thermal-maturity limit for primary thermogenic-gas generation from humic coals as determined by hydrous pyrolysis. *AAPG Bulletin*, vol. 98, p. 2581–2610.
- Liang, X. W., Wu, K., Ma, Q. F., Feng, Q. L. and Yu, J. X., 2014: Pore characteristics of different lithologies and their geological significance from the Middle Permian Gufeng Formation in the North Margin of Yangtze. *Geological Science and Technology Information*, vol. 33, p. 78–86. (in Chinese with English abstract)
- Lu, B. Q. and Qu, J. Z., 1990: Sedimentation of anoxic environments under transgression and upwelling process in Early Permian in Lower Yangtze area. *Chinese Science Bulletin*, vol. 35, p. 1193–1198.
- Ma, Q. F. and Feng, Q. L., 2012: Taxonomy and biostratigraphy of the Middle Permian radiolarian fauna from the Gufeng Formation in Luojiaba, West Hubei Province.

- Acta Micropalaeontologica Sinica*, vol. 29, p. 402–415. (in Chinese with English abstract)
- Ma, Q. F., Feng, Q. L., Carisroit, M., Danelian, T. and Zhang, N., 2016. Integrated radiolarian and conodont biostratigraphy of the Middle Permian Gufeng Formation (South China). *Comptes Rendus Palevol*, vol. 15, p. 453–459.
- Mao, Y., Ma, Q. F. and Feng, Q. L., 2013: Discovery of fish microremains in the Gufeng Formation at the Luojiaba section from Jianshi, west Hubei. *Acta Micropalaeontologica Sinica*, vol. 30, p. 175–183. (in Chinese with English abstract)
- McManus, J., Nagler, T. F., Siebert, C., Wheat, C. G. and Hammond, D. E., 2002: Oceanic molybdenum isotope fractionation: diagenesis and hydrothermal ridge-flank alteration. *Geochemistry, Geophysics, Geosystems*, vol. 3, art. no. 1078.
- Mitsumura, K. and Kamata, Y., 2009: Radiolarian faunal change across the Guadalupian–Lopingian boundary in the Sasayama section of the Tamba Belt, Southwest Japan. *News of Osaka Micropaleontologists (NOM), Special Volume*, no. 14, p. 545–559. (in Japanese with English abstract)
- Murchey, B. and Jones, D., 1992: A mid-Permian chert event: widespread deposition of biogenic siliceous sediments in coastal, island arc and oceanic basins.

Palaeogeography, Palaeoclimatology, Palaeoecology, vol. 96, p. 161–174.

Murphy, M. A. and Salvador, A., 1999: International Stratigraphic Guide — An abridged version. *Episodes*, vol. 22, p. 255–271.

Murray, R. W., 1994: Chemical criteria to identify the depositional environment of chert: general principles and applications. *Sedimentary Geology*, vol. 90, p. 213–232.

Nagai, H. and Zhu, S. Z., 1992: Permian radiolaria from the Gufeng Formation of the Nanjing area, China. *Bulletin of Nagoya University Furukawa Museum*, vol. 8, p. 1–11.

Nagai, H., Zhu, S. Z., Kametaka, M. and Wu, C., 1998: Preliminary report on Middle Permian radiolarians from the Gufeng Formation at Anmenkou, Chaohu City, Anhui Province, China. *Bulletin of Nagoya University Furukawa Museum*, vol. 14, p. 115–123.

Nakae, S., 2011: Middle and Late Permian radiolarians from the Nanjo Mountains, Fukui Prefecture, Southwest Japan. *Bulletin of the Geological Survey of Japan*, vol. 62, p. 441–453.

Nestell, G. P. and Nestell, M. K., 2010: Late Capitanian (latest Guadalupian, Middle Permian) radiolarians from the Apache Mountains, West Texas. *Micropaleontology*,

vol. 56, p. 7–68.

Nishikane, Y., Kaiho, K., Takahashi, S., Henderson, C. M., Suzuki, N. and Kanno, M.,

2011: The Guadalupian-Lopingian boundary (Permian) in a pelagic sequence from

Panthalassa recognized by integrated conodont and radiolarian biostratigraphy.

Marine Micropaleontology, vol. 78, p. 84–95.

Noble, P., Aitchison, J. C., Danelian, T., Dumitrica, P., Maletz, J., Suzuki, N., Cuvelier,

J., Caridroit, M. and O'Dogherty, L., 2017: Taxonomy of Paleozoic radiolarian

genera. *Geodiversitas*, vol. 39, p. 419–502.

Noble, P. J. and Jin, Y. X., 2010: Radiolarians from the Lamar Limestone, Guadalupe

Mountains, West Texas. *Micropaleontology*, vol. 56, p. 117–147.

Ogg, J. G., Ogg, G. M. and Gradstein, F. M., 2016: *A Concise Geologic Time Scale 2016*,

242 p. Amsterdam: Elsevier.

Petsch, S. T., Berner, R. A. and Eglinton, T. I., 2000: A field study of the chemical

weathering of ancient sedimentary organic matter. *Organic Geochemistry*, vol. 31, p.

475–487.

Prothero, D. R. and Schwab, F., 1996: *Sedimentary Geology: An Introduction to*

Sedimentary Rocks and Stratigraphy, 575 p. 7 Freeman, New York.

Saesaengseerung, D., Agematsu, S., Sashida, K. and Sardud, A., 2009: Discovery of

- Lower Permian radiolarian and conodont faunas from the bedded chert of the Chanthaburi area along the Sra Kaeo suture zone, eastern Thailand. *Paleontological Research*, vol. 13, p. 119–138.
- Saitoh, M., Isozaki, Y., Yao, J. X., Ji, Z. S., Ueno, Y. and Yoshida, N., 2013: The appearance of an oxygen-depleted condition on the Capitanian disphotic slope/basin in South China: Middle–Upper Permian stratigraphy at Chaotian in northern Sichuan. *Global and Planetary Change*, vol. 105, p. 180–192.
- Saitoh, M., Ueno, Y., Isozaki, Y., Nishizawa, M., Shozugawa, K., Kawamura, T., Yao, J. X., Ji, Z. S., Takai, K., Yoshida, N. and Matsuo, M., 2014: Isotopic evidence for water-column denitrification and sulfate reduction at the end-Guadalupian (Middle Permian). *Global and Planetary Change*, vol. 123, p. 110–120.
- Sheng, J. Z. and Wang, Y. J., 1985: Fossil Radiolaria from Kufeng Formation at Longtan, Nanjing. *Acta Palaeontologica Sinica*, vol. 24, p. 171–180. (*in Chinese with English abstract*)
- Shen, S. Z., Xie, J. F., Zhang, H. and Shi, G. R., 2009: Roadian–Wordian (Guadalupian, Middle Permian) global palaeobiogeography of brachiopods. *Global and Planetary Change*, vol. 65, p. 166–181.
- Shi, L., Feng, Q. L., Shen, J., Ito, T. and Chen, Z. Q., 2016: Proliferation of

- shallow-water radiolarians coinciding with enhanced oceanic productivity in reducing conditions during the Middle Permian, South China: evidence from the Gufeng Formation of western Hubei Province. *Palaeogeography, Palaeoclimatology, Palaeoecology*, vol. 444, p. 1–14.
- Shimmield, G. B. and Price, N. B., 1986: The behaviour of molybdenum and manganese during early sediment diagenesis—offshore Baja California, Mexico. *Marine Chemistry*, vol. 19, p. 261–280.
- Stanley, S. and Yang, X., 1994: A double mass extinction at the end of the Paleozoic Era. *Science*, vol. 166, p. 1340–1344.
- Takahashi, K. U. and Suzuki, N., 2017: Semi-open and closed system pyrolysis of Paleogene coal for evaluating the timing of hydrocarbon gas expulsion. *International Journal of Coal Geology*, vol. 178, p. 100–109.
- Takebe, M., Kametaka, M., Takayanagi, Y., Mimura, K., Sugitani, K. and Yamamoto, K., 2007: Origin and deposition of organic matter in continental chert of the Middle Permian Gufeng Formation in the northeastern Yangtze platform. *Sedimentary Geology*, vol. 201, p. 141–148.
- Trappe, J., 1998: *Phanerozoic Phosphorite Depositional Systems: A Dynamic Model for a Sedimentary Resource System*. Lecture Note in Earth Science, vol. 76. Springer,

Berlin.

Tucker, M. E., 1991: *Sedimentary Petrology: An Introduction to the Origin of Sedimentary Rocks*. 2nd ed., 260 p. Blackwell, Oxford.

van Krevelen, D. W., 1961: *Coal*, 514 p. Elsevier, Amsterdam.

Vorliceck, T. P., Kahn, M. D., Kasuya, Y. and Helz, G. R., 2004: Capture of Molybdenum in pyrite-forming sediments: role of ligand-induced reduction by polysulfides. *Geochimica et Cosmochimica Acta*, vol. 68, p. 547–556.

Wang, C. Y., 1995a: A conodont fauna from the lowermost Kufeng Formation (Permian). *Acta Micropalaeontologica Sinica*, vol. 12, p. 293–297.

Wang, Q., Ma, Z. W., Li, H. Z., Zhou, Y. Z., Liang, J., XU, L. and An, Y. F., 2008: Study on characteristics and its origin of the cherts of Gufeng Group in southern Anhui Province. *Journal of the Graduates Sun Yat-sen University (Natural Science, Medicine)*, vol. 29, p. 21–28. (in Chinese with English abstract)

Wang, R. J., 1993a: Fossil Radiolaria from Kufeng Formation of Chaohu, Anhui. *Acta Palaeontologica Sinica*, vol. 32, p. 442–457. (in Chinese with English abstract)

Wang, R. J., 1993b: Fossil radiolarians from Kufeng Formation, Hushan area, Nanjing. *Acta Micropalaeontologica Sinica*, vol. 10, p. 459–468. (in Chinese with English abstract)

- Wang, R. J., 1993c: Discovery of foraminifers in the Permian radiolarites in Jiangsu and Anhui Provinces and their depositional environment. *Journal of Tongji University*, vol. 21, p. 519–524. (in Chinese with English abstract)
- Wang, R. J., 1993d: Morphological change of the *Ruzhencevispongos uralicus* in Gufeng Formation (Lower Permian) from Jiangsu and Anhui Provinces. *Shanghai Geology*, vol. 48, p. 5–12. (in Chinese with English abstract)
- Wang, R. J., 1995b: New data of radiolarians from Gufeng Formation at Chaohu, Anhui Province. *Earth Science—Journal of China University of Geosciences*, vol. 20, p. 508–510. (in Chinese with English abstract)
- Wang, R. J., 1995c: Radiolarian fauna from Gufeng Formation (Lower Permian) in Hushan area of Nanjing, Jiangsu Province. *Scientia Geologica Sinica*, vol. 30, p. 139–146. (in Chinese with English abstract)
- Wang, R. J., Shen, G. P. and Sashida, K., 1997: Studies on radiolarian fauna from the Gufeng Formation in Anhui and Jiangsu Province, east China and its paleoenvironmental significance. *Journal of Tongji University*, vol. 25, p. 559–564. (in Chinese with English abstract)
- Wang, Y. J. and Li, J. X., 1994: Discovery of the *Follicucullus bipartitus*–*F. charveti* radiolarian assemblage zone and its geological significance. *Acta*

Micropalaeontologica Sinica, vol. 11, p. 201–212.

Wang, Y. J. and Qi, D. L., 1995: Radiolarian fauna of the Kuhfeng Formation in southern part of Jiangsu and Anhui Provinces. *Acta Micropalaeontologica Sinica*, vol. 12, p. 374–387.

Wang, Y. J. and Yang, Q., 2003: Radiolarians from the Middle Permian Kuhfeng Formation of the Renhua area, Shaoguan, Guangdong Province. *Acta Micropalaeontologica Sinica*, vol. 20, p. 333–341. (in Chinese with English abstract)

Wang, Y. J. and Yang, Q., 2007: Carboniferous–Permian radiolarian biozones of China and their paleobiogeographic implication. *Acta Micropalaeontologica Sinica*, vol. 24, p. 337–345. (in Chinese with English abstract)

Wang, Y. J. and Yang, Q., 2011: Biostratigraphy, phylogeny and paleobiogeography of Carboniferous–Permian radiolarians in South China. *Palaeoworld*, vol. 20, p. 134–145.

Wu, K., Ma, Q. F. and Feng, Q. L., 2012: Middle Permian pore characteristics and shale gas exploration significance from the Gufeng Formation in Jianshi, western Hubei. *Earth Science—Journal of China University of Geoscience*, vol. 37 p. 175–183. (in Chinese with English abstract)

- Wu, K., Ma, Q. F. and Feng, Q. L., 2015: Stratigraphic division and spatial distribution of the Middle Permian Kuhfeng Formation in the Northern Yangtze block. *Journal of stratigraphy*, vol. 39, p. 33–39. (in Chinese with English abstract)
- Xi, Y. H., 1982: Molluscas fossils from the Early Permian Gufeng Formation in Guichi Anhui Province. *Journal of Huainan Institute of Petrology (Natural Science)*, vol. 2, p. 57–70.
- Xia, B. D., Zhong, L. R., Fang, Z. and Lu. H. B., 1995: The origin of cherts of the Early Permian Gufeng Formation in the Lower Yangtze area, eastern China. *Acta Geologica Sinica*, vol. 69, p. 125–138. (in Chinese with English abstract)
- Xian, W. C. and Zhang, N., 1998: Early to Middle Permian radiolarians from the Kuhfeng Formation in southeastern Guangxi, South China. *Earth Science (Chikyu Kagaku)*, vol. 52, p. 188–202.
- Xiao, Y. F., Suzuki, N. and He, W. H., 2017: Water depths of the latest Permian (Changhsingian) radiolarians estimated from correspondence analysis. *Earth-Science Reviews*, vol. 173, p. 141–158.
- Xiao, Y. F., Suzuki, N. and He, W. H., 2018: Low-latitudinal standard Permian radiolarian biostratigraphy for multiple purposes with Unitary Association, Graphic Correlation, and Bayesian inference methods. *Earth-Science Reviews*, vol. 179, p.

168–206.

- Xie, J. C., Sun, W. D., Du, J. G., Xu, W., Wu, L. B., Yang, X. Y. and Zhou, T. F., 2013: Geochemical studies on Permian manganese deposits in Guichi, eastern China: Implications for their origin and formative environments. *Journal of Asian Earth Sciences*, vol. 74, p. 155–166.
- Xu, W., Yue, S. C., Du, J. G., Du, S. G. and Sun, C. Y., 2004: Characteristics of outcrop sequence stratigraphy of the Permian Gufeng Formation in the Guichi area, Anhui. *Journal of Stratigraphy*, vol. 28, p. 137–147. (in Chinese with English abstract)
- Yamamoto, K., 1987: Geochemical characteristics and depositional environments of cherts and associated rocks in the Franciscan and Shimanto terranes. *Sedimentary Geology*, vol. 52, p. 65–108.
- Yang, S. Y. and Yao, J., 2008: Geochemistry and origin of siliceous rocks from the Gufeng Formation of Middle Permian in the Pingdingshan area, Chaohu region, Anhui Province. *Geological Journals of China Universities*, vol. 14, p. 39–48. (in Chinese with English abstract)
- Yang, Y. Q. and Feng, Z. Z., 1997: Formation and significance of the bedded siliceous rocks of the Lower Permian in South China. *Acta Petrologica Sinica*, vol. 13, p. 111–119. (in Chinese with English abstract)

- Yao, A., Kuwahara, K., Yao, J. X., Ji, Z. S. and Li, J. X., 2007: Permian radiolarians from the so-called Gufeng Formation of the Laibin-Liuzhou area, Guangxi, China. *Journal of Geosciences, Osaka City University*, vol. 50, p. 67–81.
- Yao, A., Yu, J. and An, T. X., 1993: Late Paleozoic radiolarians from the Guizhou and Guangxi areas, China. *Journal of Geosciences, Osaka City University*, vol. 36, p. 1–13.
- Yao, X., Zhou, Y. Q. and Hinnov, L. A., 2015: Astronomical forcing of a Middle Permian chert sequence in Chaohu, South China. *Earth and Planetary Science Letters*, vol. 422, p. 206–221.
- Ye, L. F. and Li, J., 1924: Coalfield geology of Jixian, Anhui*. *Geological Report**, vol. 6, p. 25–50. (*in Chinese*) *translated by the authors
- Zeng, P. S., Yang, Z. S., Meng, Y. F., Pei, R. F., Hou, Z. Q., Xu, W. Y., Wang, X. C. and Tian, S. H., 2004: Petrogenesis and significance of cherts in Tongling mineralization cluster area, Anhui. *Geological Review*, vol. 50, p. 153–161. (*in Chinese with English abstract*)
- Zhang, B. L., Yao, S. P., Hu, W. X., Ding, H., Liu, B. and Ren, Y. L., in press: Development of a high-productivity and anoxic-euxinic condition during the late Guadalupian in the Lower Yangtze region: Implications for the mid-Capitanian

extinction event. *Palaeogeography, Palaeoclimatology, Palaeoecology*.

<https://doi.org/10.1016/j.palaeo.2018.01.021>

Zhang, L., Ito, T., Feng, Q. L., Caridroit, M., and Danelian, T., 2014: Phylogenetic model of *Follicucullus*-lineages (Albaillellaria, Radiolaria) based on high resolution biostratigraphy of the Permian Bancheng Formation, Guangxi, South China. *Journal of Micropalaeontology*, vol. 33, p. 179–192.

Zhang, N., Henderson, C. M., Xia, W. C., Wang, G. Q. and Shang, H. J., 2010: Conodonts and radiolarian through the Cisuralian–Guadalupian boundary from the Pingxiang and Dachongling sections, Guangxi region, South China. *Alcheringa: An Australian Journal of Palaeontology*, vol. 34, p. 135–160.

Zhao, J. K. and Zheng, Z. G., 1977: The Permian ammonoids from Zhejiang and Jiangxi. *Acta Palaeontologica Sinica*, vol. 16, p. 217–252. (in Chinese with English abstract)

Zhou, X. P., He, Y. B., Du, H. Q. and Li, H., 2009: Geochemical characteristics and origin of the Permian siliceous rocks in Xuanhan region of Sichuan Province. *Journal of Paleobiogeography*, vol. 11, p. 670–680. (in Chinese with English abstract)

Zhu, S. Z., Nagai, H., Kametaka, M. and Wu, C. Y., 1999: Nectic chert, a very peculiar

type of siliceous rock. *Geological Journals of China Universities*, vol. 5, p. 345–350.

(in Chinese with English abstract)

Zhu, T. X., 1989: Sedimentological features and the genesis of the lower Permian nodular and thin-bedded siliceous rocks in southern Anhui. *Sedimentary Facies and Palaeogeography*, vol. 43, p. 1–8. (in Chinese with English abstract)

Zhu, Z. Y., Jiang, S. Y., Liu, G. X. and Zhao, K. D., 2013: Precise dating of the Middle Permian: zircon U–Pb geochronology from volcanic ash beds in the basal Gufeng Formation, Yangtze region, South China. *Gondwana Research*, vol. 23, p. 1599–1606.

Ziegler, A. M., Hulver, H. L. and Rowley, D. B., 1997: Permian world topography and climate. In, Martini, I. P. ed., *Late Glacial and Postglacial Environmental Changes—Quaternary, Carboniferous–Permian and Proterozoic*, p. 111–146. Oxford University Press, New York.

Figure captions

Figure 1. Guadalupian paleogeographic map, modified from Ziegler *et al.* (1997).

Figure 2. Original and emended definitions of the Gufeng Formation. Note that the information by Ye and Li (1924) is re-quoted from Bureau of Geology and Mineral Resources of Anhui Province (1997). English of the definitions are translated by the authors.

Figure 3. Distributions of the Gufeng Formation with the major localities of sections and the radiolarian localities from previous studies. The distributions are based on Wu *et al.* (2015). Detailed location of the Lengshuixi section in Shizhu was not shown by Bureau of Geology and Mineral Resources of Sichuan Province (1997); therefore, the location is not presented.

Figure 4. Lithostratigraphy of the stratotype section of the Gufeng Formation in Anhui Province and the hypostratotype sections in the Hubei, Jiangsu, Zhejiang, Hunan, and Guangdong provinces with major sections. White triangles with alphabets indicate occurrence horizons of each photograph of Figure 5.

Figure 5. Field occurrences of the Gufeng Formation. **A, B**, black bedded cherts in the Liuhuang section in Chaohu, Anhui; **C**, overturned basal mudstone and



stratigraphically overlying limestone of the Maokou Formation in the Liuhuang section; **D**, black bedded cherts of the Maocaojie section in Jianshi, Hubei; **E**, mudstone including lenticular limestone in the Luojiaba section in Jianshi; **F**, carbonaceous mudstone and stratigraphically overlying limestone of the Wuxue Formation in the Maocaojie section. The occurrence horizons of each correspond to the white triangles in Figure 4.

Figure 6. Summarized radiolarian zones of the Gufeng Formation in each province in previous studies. Note that the *Follicucullus monacanthus* Zone is changed to *Pseudoalbaillella monacantha* Zone according to the taxonomic reconsideration by Ito *et al.* (2015). *F.*, *Follicucullus*; *R.*, *Ruzhencevispongos*; *Ps.*, *Pseudoalbaillella*; *A.*, *Albaillella*.


Figure 7. SEM images of major remarkable radiolarians from the Gufeng Formation (reprinted from Ito *et al.*, 2013a). **A, B**, *Longtanella* cf. *zhengpanshanensis* Sheng and Wang; **C**, *Longtanella* sp.; **D, E**, *Ruzhencevispongos uralicus* Kozur; **F**, *Hegleria* cf. *mammilla* (Sheng and Wang).


Figure 8. Sketch of a holotype specimen of *Longtanella zhengpanshanensis* Sheng and Wang, a type species of genus *Longtanella* Sheng and Wang (drawn from Sheng and Wang, 1985).


Figure 9. Schematic model showing vertical changes of lithology, radiolarian fauna, and redox condition. Numerical ages with white arrows indicate zircon U–Pb ages of 272.0 ± 5.5 Ma (MSWD = 2.6) and 271.5 ± 3.3 Ma (MSWD = 1.7) separated from volcanic ash beds within the basal shale strata of the Gufeng Formation in Chaohu area, Anhui by Zhu *et al.* (2013). Lop., Lopingian; Cis., Cisuralian; *F.*, *Follicucullus*; *R.*, *Ruzhencevisponus*; *Ps.*, *Pseudoalbaillella*; MK, Maokou Formation; WX, Wuxue Formation; YP, Yinpin Formation; UU, Upper Unit.




	Author	Name	Definition
Original definiton	叶良辅·李捷（1924）	孤峰镇石灰岩	为黑色灰质頁岩及硅质甚富之灰岩等，间干栖霞灰岩及龙潭煤系间，底部之灰质黑色页岩含化石极丰，其最著者为菊石及腕足类。
	Ye and Li (1924)	Gufeng Zheng Limestone	It is characterized by black carbonate shales and silica-rich carbonates, which are lithostratigraphically located between the Qixia and Longtan formations. Basal black shales include abundant fossils represented by ammonite and brachiopod.
Emended definiton	安徽省地质矿产局（1997）	孤峰組	指栖霞組与龙潭組之间的薄层状硅质岩、硅质页岩、粉砂质泥岩、碳质页岩、锰质页岩地层，下部页岩含锰及磷結核。 产菊石、腕足类、双壳类及放射虫等多门类化石。 底与栖霞組平行不整合接触，以厚层灰岩消失、含锰页岩出現为界； 頂与龙潭組整合接触，以薄层硅质岩消失、页岩出現为界。
	Bureau of Geology and Mineral Resources of Anhui Province (1997)	Gufeng Formation	It is characterized by black or gray-yellow thin-bedded siliceous rocks, siliceous shales, siltstones, carbonous shales, and manganese shales. The Gufeng Formation overlies the Qixia Formation and the underlies the Longtan Formation. Basal shales include manganese and phosphate nodules. Several fossils (e.g., ammonoids, brachiopods, bivalves, and radiolarians) occur from it.





 Distribution of the Gufeng Formation

 Radiolarian locality with columnar section in Figure 4

 Radiolarian locality

 Hypostratotype sections established by the bureaus of geology and mineral resources in each province







A



B




limestone
(Maokou Fm.)

mudstone

C



D



lenticular limestone

mudstone

lenticular limestone

E



limestone
(Wuxue Fm.)


carbonaceous
mudstone

F




Age		Southwest Japan (Ishiga, 1986, 1990; Kuwahara <i>et al.</i> , 1998)	South China (Wang and Yang, 2007, 2011)	North margin			South margin	
				Gufeng Formation			“Gufeng Formation”	
				Upper Yangtze Region (Shichuan) (Kuwahara <i>et al.</i> , 2007c, 2008b)	Middle Yangtze Region (Hubei) (Feng and Zhong, 1994; Kuwahara <i>et al.</i> , 2007a, 2008a; Ma and Feng, 2012; Ma <i>et al.</i> , 2016; Shi <i>et al.</i> , 2016)	Lower Yangtze Region (Anhui and Jiansu) (e.g., Sheng andWang, 1985; Wang and Qi, 1995; Wang <i>et al.</i> , 1997; He <i>et al.</i> , 1999; Kuwahara <i>et al.</i> , 2007b; Kametaka <i>et al.</i> , 2009; Ito <i>et al.</i> , 2013a)	Guangxi (Yao <i>et al.</i> , 1993; Wang and Li, 1994; Xian and Zhang, 1998; Kuwahara <i>et al.</i> , 2004)	Guangdong (Wang and Yang, 2003)
Permian	Lopingian (part)	<i>Follicucullus charveti</i> - <i>Albaillella yamakitai</i> Assemblage Zone	<i>F. bipartitus</i> - <i>F. charveti</i> - <i>F. orthogonus</i> Assemblage Zone				<i>F. charveti</i> - <i>A. yamakitai</i> Assemblage Zone	
	Guadalupian	<i>F. ventricosus</i> - <i>F. scholasticus</i> Assemblage Zone	<i>F. scholasticus</i> - <i>F. ventricosus</i> Assemblage Zone	<i>F. scholasticus</i> - <i>R. uralicus</i> Assemblage Zone	<i>F. scholasticus</i> Assemblage Zone	<i>F. scholasticus</i> - <i>R. uralicus</i> Assemblage Zone	<i>F. ventricosus</i> - <i>F. scholasticus</i> Assemblage Zone	
		<i>Ps. monacantha</i> Range Zone	<i>Ps. monacantha</i> Range-Zone		<i>Ps. monacantha</i> Assemblage Zone	<i>Ps. monacantha</i> Assemblage Zone		
		<i>Ps. globosa</i> Assemblage Zone	<i>Ps. globosa</i> Range Zone	<i>Ps. longtanensis</i> - <i>Ps. fusiformis</i> Assemblage Zone	<i>Ps. globosa</i> -Assemblage Zone	<i>Ps. longtanensis</i> - <i>P. fusiformis</i> Assemblage Zone		<i>Ps. fusiformis</i> - <i>Ps. globosa</i> Assemblage Zone
	Cisuralian (part)	<i>Ps. longtanensis</i> Assemblage Zone	<i>Ps. Ishigai</i> Abundance zone				<i>P. banchengensis</i> Assemblage Zone	
		<i>A. sinuata</i> Range Zone	<i>A. sinuata</i> Abundance Zone <i>A. xiaodongensis</i> Range Zone				<i>Albaillella sinuata</i> Assemblage Zone	
		<i>Ps. scalprata</i> m. <i>rhombothoracata</i> Assemblage Zone	<i>Ps. rhombothoracata</i> Assemblage Zone				<i>P. scalprata</i> Assemblage Zone	



100 μm (A)




100 μm (B, C)



100 μm (D-F)

100 μm



Epoch / Age (Ogg <i>et al.</i> , 2016)			Radiolarian occurrence		Middle Yangtze region Maocaojie section (Shi <i>et al.</i> , 2016)		Lower Yangtze region Anmenkou section (Kametaka <i>et al.</i> , 2005, 2009)					
			Assemblage zone	Faunal change	Lithostratigraphy	Redox condition	Lithostratigraphy	Redox condition				
<div>(Ma)</div> <div>260</div> <div>265</div> <div>270</div> <div>271.5±3.3▽ 272.0±5.5▽ Zhu <i>et al.</i> (2013)</div>	Permian	Lop.	Wuchiapingian 259.8			WX	<div>carbonate</div>		YP	<div>clastics</div>		
		Guadalupian			no radiolarian occurrence		UU	<div>muddy rock</div>	suboxic		<div>muddy rock</div>	
			Capitanian	<i>F. scholasticus</i> – <i>R. uralicus</i>	spherical radiolaria-dominant ↑	Gufeng Formation	Middle Unit	<div>muddy rock with carbonate</div>	anoxic	SRM	<div>chert</div>	anoxic
			Wordian	<i>Ps. monacantha</i>								
			Roadian	<i>Ps. globosa</i>			Lower Unit	<div>chert</div>				
		Cis.	Kungrian			MK	<div>carbonate</div>		MK	<div>carbonate</div>		
									PNNM	<div>muddy rock with nodules</div>	suboxic aerobic	