

Paleontological Research



Papers in Press

“Papers in Press” includes peer-reviewed, accepted manuscripts of research articles, reviews, and short notes to be published in *Paleontological Research*. They have not yet been copy edited and/or formatted in the publication style of *Paleontological Research*. As soon as they are printed, they will be removed from this website. Please note they can be cited using the year of online publication and the DOI, as follows:

Humblet, M. and Iryu, Y. 2014: Pleistocene coral assemblages on Irabu-jima, South Ryukyu Islands, Japan. *Paleontological Research*,
doi: 10.2517/2014PR020.

Middle and Late Jurassic radiolarians from Nadanhada terrane of eastern Heilongjiang Province, northeastern China

GANG LI¹, ATSUSHI MATSUOKA², QUN YANG¹ AND JINGENG SHA¹

¹State Key Laboratory of Palaeobiology and Stratigraphy, Center for Excellence in Life and Palaeoenvironment, Nanjing Institute of Geology and Palaeontology, Chinese Academy of Sciences, Nanjing 210008, China. (*gangli@nigpas.ac.cn*)

²Faculty of Science, Department of Geology, Niigata University, Niigata 950-2181, Japan.

Abstract. The Nadanhada terrane of northeastern Heilongjiang, northeastern China is composed of Jurassic accretionary complexes (i.e., the Yuejinshan Complex in the west, the Raohe Complex in the east) and succedent cover beds. In this paper we report two radiolarian assemblages recognized from four black claystone samples collected from the Dalingqiao Formation in the Dajiashan area of the Raohe Complex. The *Striatojaponocapsa synconexa*-“*Tricolocapsa*” *tetragona* assemblage of one sample consists of eleven species in eight genera, which indicates a middle Bathonian (Middle Jurassic) age, i.e., correlative with the uppermost *Striatojaponocapsa*

plicarum Zone to the lower *Striatojaponocapsa conexa* Zone. The other three samples contain 34 species in 25 genera, which indicate a late Oxfordian–early Tithonian (Late Jurassic) age. The discovery of the Late Jurassic radiolarian assemblage indicates that the Raohe Accretionary Complex was formed during the Late Jurassic. The succedent late Tithonian–early Valanginian *Buchia* fauna-bearing Dong’anzen Formation may be the earliest cover beds overlying the Raohe Accretionary Complex. Then, during the Early Cretaceous, the Nadanhada terrane received the deposition of the *Aucellina* bivalve fauna-bearing Dajiashan Group. This indicates paleogeographic differentiation between the Nadanhada terrane and the Tamba-Mino-Ashio terrane.

Key words: eastern Heilongjiang, Middle–Late Jurassic, Nadanhada terrane, northeastern China, radiolarian biostratigraphy

Introduction

The early geological study on the Nadanhada terrane of northeastern China began in 1930s by Japanese geologists (Yabe and Ohki, 1957). In 1957 Chinese and Russian scientists made a joint geological research in the Ussuri River region, and confirmed the occurrence of Late Triassic, Early and Middle Jurassic rocks based on the fossil evidence (Wang, 1959), and they came to the conclusion that the Mesozoic mobile belt developed in a geosyncline in the northeastern continental margin of East Asia. Based on fossil records of Late Paleozoic fusulinids the Nadanhada terrane was re-interpreted as a Paleozoic geosyncline that has extended from the northeast

Sikhote-Alin region (Li *et al.*, 1979). But the sporadically and irregularly distributed rocks yield various age fossils, such as Carboniferous and Permian fusulinids in limestone, Triassic conodonts in bedded chert (Wang *et al.*, 1986; Buryi, 1996), Triassic, Early and Middle Jurassic radiolarians in bedded chert and siliceous shale (Kojima and Mizutani, 1987; Yang *et al.*, 1993) and Late Jurassic to Early Cretaceous *Buchia* bivalves in claystone of the Dong'anzen Formation (Sun *et al.*, 1989; Sha and Fürsich, 1993; Sha *et al.*, 1994, 2008, 2009). These research results strongly suggest the melange geological characters of the Nadanhada terrane, which together with the Mino and Western Sikhote-Alin terranes formed a single superterrane before the opening of the Japan Sea (Kojima, 1989). This interpretation would be very helpful to understand and decipher the paleo-Pacific-subduction history (Wu *et al.*, 2007; Zhou *et al.*, 2009, 2014; Kojima *et al.*, 2016).

Recent studies demonstrated that the Dajiashan Group, cropping out in the Dajiashan area, southern Nadanhada terrane, yields a middle-late Early Cretaceous *Aucellina* bivalve fauna (Sha, 1990, 2002, 2007). Furthermore, the purported Early Jurassic ammonites of the Dajiashan Group (Li, 1996) are also similar to those of a *Pseudohaploceras* ammonite fauna from the Lower Cretaceous Longzhaogou Group (Futakami *et al.*, 1995; Li and Bengtson, 2018). All these sparked an interest in restudying the paleontology and biostratigraphy of the Dajiashan Group and its underlying deposits.

Well-preserved Middle-Late Jurassic radiolarian faunas are encountered in four samples from the black claystone of the Dalingqiao Formation, which was originally

assigned to the Upper Triassic–Lower Jurassic by the Geological Survey of Heilongjiang Province of China (Bureau of Geology and Mineral Resources of Heilongjiang Province, 1993). Detailed taxonomical and biostratigraphical study of the recovered radiolarian faunas of the Dalingqiao Formation would help us to understand the geology of the Nadanhada terrane.

Material and methods

The fieldwork was carried out by the first author in August and September of 2002, and concerned the Dajiashan area, north Hulin. The section P1 was investigated (Figure 1). Some ammonites and bivalves were encountered in the Lower Cretaceous Dajiashan Formation (Figure 2). Although many samples for microfossil analysis were collected, relatively abundant and better-preserved radiolarians were recovered in four samples (P1-08, 09, 012, 013) (Figure 3) from the Upper Jurassic Dalingqiao Formation, which underlies the Dajiashan Group. Although the four samples were collected in the order of from upper to lower horizons (as is shown in Figure 3), the poor outcrop condition can not clearly show their real relative horizons.

In order to extract the radiolarian tests, the samples were treated after the method initiated by Dumitrica (1970) and described later by Pessagno and Newport (1972). The sample rocks were first crushed into centimeter-sized fragments and treated with 5% hydrofluoric acid (HF) for about 20 hours. The samples were washed in a two-sieved set, the lower one with a mesh diameter of 63 μm , the upper one with a mesh

diameter of 315 μm . The fine residues in the lower sieve were dried; the coarser residues in the upper sieve were put back into the plastic beaker, and processed with hydrofluoric acid once more. The radiolarians were then picked from the dry residues with a brush pen bearing two to three hairs under a stereomicroscope and preliminary determined. The better preserved specimens were mounted on stubs and examined and photographed under the Scanning Electron Microscope LEO 1530 VP (the State Key Laboratory of Palaeobiology and Stratigraphy) for more precise determinations.

The section P1 is poorly exposed, and located in the forest in north of Hulin County, about 12 km southeast of Wulindong, Raohe County. The Dajiashan Group is well developed. The section was measured, by the help of test trenches along the top of hills, by the Regional Geological Survey of the Bureau of Geology and Mineral Resources of Heilongjiang Province.

In the P1 section the Dajiashan Group is subdivided into three formations: i.e., in ascending order, the Nandatashan, Dajiashan and Baiheshan formations (Figure 2). The Nandatashan Formation is a volcanic-clastic sequence with a thickness of 80 m. The Dajiashan Formation (263 m thick) consists of dark gray claystone, fine-grained sandstone and siltstone, yielding ammonites. The Baiheshan Formation (243 m thick) consists of black claystone intercalating fine-grained sandstone and siltstone, yielding abundant bivalves. The here described Middle and Late Jurassic radiolarians are from the strata classified as the Dalingqiao Formation in the P1 section.

Radiolarian assemblages and their age assignments

Well-preserved Middle–Late Jurassic radiolarians are encountered in four samples (P1-013, 012, 09, 08) (Table 1) from black claystone in the P1 section of the Dalingqiao Formation, which was originally assigned to the Upper Triassic–Lower Jurassic by the Heilongjiang Geological Survey (Bureau of Geology and Mineral Resources of Heilongjiang Province, 1993). These radiolarians encompass 43 species belonging to 31 genera. Two different-aged radiolarian assemblages have been recognized (Figure 4). The sample P1-09 yields a late Middle Jurassic assemblage, which could be assigned to the uppermost part of the *Striatojaponocapsa plicarum* Zone to the basal part of the *Striatojaponocapsa conexa* Zone (Matsuoka, 1995). The other three samples (P1-08, 012, 013) yield a Late Jurassic assemblage, which could be most probably assigned to the Kimmeridgian *Hsuum maxwelli* Zone (Matsuoka, 1995).

1. *Striatojaponocapsa synconexa*-“*Tricolocapsa*” *tetragona* assemblage

This assemblage consists of eleven species belonging to eight genera (except for Cryptothracic Nassellaria), they are: *Archaeocenosphaera* sp., *Caneta?* sp. 1, “*Stichocapsa*” *convexa* Yao, 1979, “*Stichocapsa*” *robusta* Matsuoka, 1984, “*Stichocapsa*” sp., *Striatojaponocapsa synconexa* (O’Dogherty *et al.*, 2005), *Tethysetta dhimenaensis* (Baumgartner, 1984), *Transhsuum* sp., “*Tricolocapsa*” sp., “*Tricolocapsa*” *tetragona* Matsuoka, 1983 and *Unuma gordus* Hull, 1997. Among

them some species have geologically long ranges. Such as “*Stichocapsa*” *convexa* has a geological range of early Callovian–late Oxfordian according to Aita (1987), or early Aalanian–early Tithonian according to Baumgartner *et al.* (1995). “*Stichocapsa*” *robusta* ranges from latest Bajocian to early Callovian (Baumgartner *et al.*, 1995), or from early Callovian to late Oxfordian (Aita, 1987). *Tethysetta dhimenaensis* has a geological range of early Bajocian–early Oxfordian. *Striatojaponocapsa synconexa* has a geological range of UAZone 4–5 of Baumgartner *et al.* (1995). While “*Tricolocapsa*” *tetragona* has a restricted geological range, i.e., UAZone 5 of Baumgartner *et al.* (1995) or the uppermost part of the “*Striatojaponocapsa*” *plicarum* Zone to the basal part of the *Striatojaponocapsa conexa* Zone of Matsuoka (1995).

2. *Cinguloturris carpatica* assemblage

This assemblage consists of 34 species belonging to 25 genera, they are: *Arcanicapsa funatoensis* (Aita, 1987), *Archaeodictyomitra rigida* Pessagno, 1977a, *Archaeodictyomitra* sp., *Archaeospongoprunum* sp. 1, *Archaeospongoprunum* sp. 2, *Caneta?* sp. 1, *Caneta* sp. 2, *Emiluvia premyogii* Baumgartner, 1984, *Cinguloturris fusiforma* Hori, 1999, *Cinguloturris carpatica* Dumitrica in Dumitrica and Mello, 1982, *Crococapsa hexagona* (Hori, 1999), *Eucyrtidiellum ptyctum* (Riedel and Sanfilippo, 1974), *Eucyrtidiellum unumaense* (Yao, 1979), *Gongylothorax favosus* Dumitrica, 1970, *Loopus* sp. 1, *Loopus* sp. 2, *Loopus?* sp. 3, *Olanda?* sp., *Orbiculiforma* sp., *Paronaella* sp. 1, *Paronaella* sp. 2, *Plegmosphaera?* spp., *Praeconocaryomma?* sp., *Protunuma japonicus* Matsuoka and Yao, 1985, *Pseudoxitus?*

sp., *Ristola* sp., *Solenotryma ichikawai* Matsuoka and Yao, 1985, *Tethysetta dhimenaensis* (Baumgartner, 1984), *Theocapsomella himedaruma* (Aita, 1987), *Triactoma* sp., *Unuma gordus* Hull, 1997, *Williriedellum marcucciae* (Cortese, 1993), *Zhamoidellum ovum* Dumitrica, 1970, and *Zhamoidellum* sp.

Among them, *Arcanicapsa funatoensis* ranges from early Callovian to middle Kimmeridgian (Aita, 1987). *Archaeodictyomitra rigida* ranges from middle Oxfordian–early Kimmeridgian to Berriasian or younger (Hull, 1997). *Cinguloturris fusiformis* was reported in the Tithonian of Japan and the Far East of Russia (Hori, 1999). *Cinguloturris carpatica* is very common in the sample P1–08, it ranges from the *Kilinora spiralis* Zone to *Loopus primitivus* Zone in Japan, and of an age of late Callovian to late Tithonian (Matsuoka and Yao, 1986; Matsuoka, 1995). *Crococapsa hexagona* was described from the Tithonian of Japan (Hori, 1999). *Zhamoidellum ovum* Dumitrica ranges from middle Oxfordian to early Tithonian according to Baumgartner *et al.* (1995). With respect to the co-occurrence of these important species, the radiolarian assemblage is most probably assigned to the *Hsuum maxwelli* Zone (Figure 4), including the upper part of the *Kilinora spiralis* Zone and the lower part of the *Loopus primitivus* Zone (Upper Oxfordian–Lower Tithonian) (Matsuoka, 1995), although the zonal radiolarian species are not discovered in the three samples.

Discussion on tectonics and paleogeography

The Nadanhada terrane (Mizutani *et al.*, 1986; Kojima, 1989; Zhang *et al.*, 1997),

located in northeastern Heilongjiang, northeastern China, contains two accretionary complexes related to paleo-Pacific plate subduction and accretion (Kojima and Kametaka, 2000; Wilde and Zhou, 2015; Kojima *et al.*, 2016): the Yuejinshan Complex in the west and the Raohe Complex in the east (Zhou *et al.*, 2014). The Yuejinshan Complex (consisting of meta-clastic rocks and metamafic-ultramafic rocks) is the first stage accretionary complex that accreted during the late Early Jurassic and led to the collision of the Jiamusi Massif westwards with the Central Asian Orogenic Belt (Wu *et al.*, 2007). The Raohe Complex, the main part of the Nadanhada terrane, is composed of Carboniferous to Permian limestone (Li *et al.*, 1979), Middle Triassic–Early Jurassic bedded chert, Middle Jurassic siliceous shale (Mizutani *et al.*, 1986; Kojima, 1989; Yang *et al.*, 1993), mafic-ultramafic rocks, which as olistoliths are inbedded in clastic “matrix”. Although there is no fossil evidence, the geological age of the clastic “matrix” was assigned to Late Jurassic or Early Cretaceous by Kojima (1989). Now, we have found four samples (P1-08, 09, 012, 013) yielding two radiolarian assemblages from the Dalingqiao Formation in the P1 section in the Dajiashan area (Figures 3, 4). Three samples (P1-08, 012, 013) contain a Late Jurassic radiolarian assemblage, and one sample (P1-09) yields a Middle Jurassic radiolarian assemblage (Figure 3). As the four radiolarian samples are not in normal stratigraphic order, we interpret that the Dalingqiao Formation as the clastic “matrix” of the accretionary melange, and the Late Jurassic radiolarian assemblage would precisely constrain the Late Jurassic accretion of the main part of the Nadanhada terrane. After the accretion event, the succedent late Tithonian–early Valanginian *Buchia*

fauna-bearing Dong'anzen Formation would be the earliest cover beds overlying the Raohe Complex. The Lower Cretaceous Dajiashan Group, containing an *Aucellina* bivalve fauna (Sha, 2007; Sha *et al.*, 2009), is contemporaneous with the Longzhaogou Group and the Jixi Group of the Jiamusi Massif (Li and Yang, 2003; Li and Yu, 2004; Li and Matsuoka, 2015; Li and Bengtson, 2018). Although Late Jurassic to Early Cretaceous radiolarian faunas were reported from the Tamba-Mino-Ashio terrane in southwestern Japan, the late Tithonian to early Valanginian *Buchia* fauna is not reported there (Sha *et al.*, 1994). And the Early Cretaceous *Aucellina* fauna is not recovered in Japan (Sha *et al.*, 1994). These faunal differences may indicate paleogeographic differentiation between the Nadanhada terrane and the Tamba-Mino-Ashio terrane.

Paleontological description

For species, reference to the author, the first illustration, and the currently adopted species concept are given. Genera and species are arranged in alphabetical order within orders Spumellaria and Nassellaria, respectively.

Order Spumellaria Ehrenberg, 1875, emend. De Wever *et al.*, 2001

Genus *Archaeocenosphaera* Pessagno and Yang, 1989

Type species.—*Archaeocenosphaera ruesti* Pessagno and Yang in Pessagno *et al.*, 1989.

Archaeocenosphaera sp.

Figures 5.17, 8.1

Material.—Specimen P1-8-29 is from the sample P1-08; specimen P1-09-25 is from the sample P1-09.

Dimension (in μm).—Diameter of cortical shell (Max./Min.): 256/241 (P1-8-29), 156/153 (P1-09-25).

Remarks.—Only two specimens are encountered, the cortical test possesses denser pore frames than *Archaeocenosphaera ruesti* Pessagno and Yang in Pessagno *et al.*, 1989, and it is easily differentiated between them. The Chinese specimens possess massive nodes in the lattices of pore frames, and differ from *A. laseekensis* Pessagno and Yang in Pessagno *et al.*, 1989.

Range and occurrence.—Middle and Upper Jurassic. Nanhada.

Genus *Archaeospongoprunum* Pessagno, 1973

Type species.—*Archaeospongoprunum venadoense* Pessagno, 1973.

Archaeospongoprunum sp. 1

Figure 8.3

Material.—Specimen P1-08-24 is from the sample P1-08.

Dimensions (in μm).—Length of spine: 196; length of test: 237; width of test: 193.

Remarks. —Test ellipsoidal, with two triradiate polar spines.

Range and occurrence.—Upper Jurassic. Nadanhada.

Archaeospongoprunum sp. 2

Figure 8.4

Material.—Specimen 108x2-08 is from the sample P1-08.

Dimensions (in μm).—Length of test: 196; width of test: 96.

Remarks.—Test cylindrical, surface ornamented with polygonal pore frames, two polar spines triradiate in axial section.

Range and occurrence.—Upper Jurassic. Nadanhada.

Genus *Emiluvia* Foreman, 1973; emend. Pessagno, 1977a

Type species.—*Emiluvia chica* Foreman, 1973.

Emiluvia premyogii Baumgartner, 1984

Figures 8.5, 8.6

Emiluvia premyogii Baumgartner, 1984, p. 762, pl. 3, figs. 6, 8, 9; Hull, 1997, p. 68,
pl. 27, figs. 1, 2, 21.

Material.—Specimens 108x2-20, P1-08-28 are from the sample P1-08.

Dimension (in μm).—Length of long spine, length of short spine, between base of spines, between concave sides: 143, 97, 130, 99 (108x2-20); 262, 209, 184, 134 (P1-08-28).

Remarks.—The small specimens are preserved with broken spines. About 6 pairs of nodes between opposed spines. Center of cross forms a raised polygonal structure often with a center node. Additional lateral meshwork has no significant node.

Occurrence.—UAZones 3–10, early–middle Bajocian to late Oxfordian–early Kimmeridgian. Blake Bahama Basin, California, Carpathians, Italy, Japan, Nadanhada, Yugoslavia.

Genus *Orbiculiforma* Pessagno, 1973

Type species.—*Orbiculiforma quadrata* Pessagno, 1973

Orbiculiforma sp.

Figure 7.3

Material.—Specimen 13-7 is from the sample P1-013.

Dimensions (in μm).—Test diameter: 240; diameter of central cavity: 120.

Remarks.—Only one specimen encountered, it is similar to *Orbiculiforma lowreyensis* Pessagno, 1977a in having sloping sides and angled periphery. The former differs from the later in having smaller test size and relative larger ratio of central cavity to

test diameter.

Range and occurrence.—Middle Jurassic. Nadanhada.

Genus *Paronaella* Pessagno, 1971, emend. Baumgartner, 1980

Type species.—*Paronaella solanoensis* Pessagno, 1971.

Paronaella sp. 1

Figure 8.7

Material.—Specimen 108x2-34 is from the sample P1-08.

Dimensions (in μm).—Length of rays: AX (A axis), 200; BX (B axis), 219; CX (C axis), 238; narrowest width of rays: AX, 86; BX, 86; CX, 76; width of ray tips: AX, 133; BX, 133; CX, 124.

Remarks.—The present specimen is preserved with faint polygonal pore frames in the surface of rays and the central part, the spines at the ray tips are broken, with respect to the test outline and the dimensions, it is closely related to *Paronaella mulleri* Pessagno, 1977a. But the lack of information of the inner structure makes further comparison difficult.

Range and occurrence.—Upper Jurassic. Nadanhada.

Paronaella sp. 2

Figure 8.8

Material.—Specimen P1-08-26 is from the sample P1-08.

Dimensions (in μm).—Length of rays: AX (A axis), ?; BX (B axis), 216; CX (C axis), 168; width of base of rays: AX, 112; BX, 112; CX, 104; length of ray spines: AX, ?; BX, ?; CX, 24.

Remarks.—The specimen is closely related to *Paronaella pygmaea* Baumgartner, 1980 except the large size of the present specimen. The poor preservation and the lack of information on the inner structure, make further identification difficult.

Range and occurrence.—Upper Jurassic. Nadhanhada.

Genus *Plegmosphaera* Haeckel, 1881

Type species.—*Plegmosphaera maxima* Haeckel, 1887.

Plegmosphaera? spp.

Figures 6.1, 6.2

Material.—Specimen P1-8-16 is from the sample P1-08; specimen P1-9-10 is from the sample P1-09; specimens P1-12-1, 6, 7, 29, 31, 32, 33 are from the sample P1-012; specimen P1-13-2 is from the sample P1-013

Dimensions (in μm).—Diameter of cortical shell (Max./Min.): 144/135 (P1-8-16), 235/209 (P1-9-10), 291/257 (P1-12-1), 224/223 (P1-12-6), 219/206 (P1-12-7), 230/222 (P1-12-29), 249/238 (P1-12-31), 145/132 (P1-12-32), 148/128 (P1-12-33),

173/168 (P1-13-2).

Remarks.—The Chinese specimens are spherical or ellipsoidal spongy spumellarians, they should be differentiated into different species, but the present specimens are not preserved with inner structure, and are preliminarily included in the genus.

Range and occurrence.—Late Callovian to Berriasian. Antarctica, Italy, Nanhada, Oman.

Genus *Praeconocaryomma* Pessagno, 1976

Type species.—*Praeconocaryomma universa* Pessagno, 1976.

Praeconocaryomma? sp.

Figures 7.1, 7.2, 8.2

Material.—Specimen P1-08-17 is from the sample P1-08; Specimens P1-13-5, 12, 14 are from the sample P1-013.

Description.—Cortical test spherical, having numerous mammae in lower relief, and having triangular meshwork in the area between mammae.

Dimension (in μm).—Diameter of cortical test 208 (P1-08-17), 260/247 (P1-13-5), 253/207 (P1-13-12), 256/249 (P1-13-14).

Remarks.—The Chinese specimen is similar to *Praeconocaryomma mamillaria* (Rüst) by sharing triangular meshwork in the area between small mammae in lower relief, but the present specimen is poorly preserved, and this hampers further comparison.

Range and occurrence.—Upper Jurassic. Nadanhada.

Genus *Triactoma* Rüst, 1885

Type species.—*Triactoma tithonianum* Rüst, 1885, subsequent designation by Campbell, 1954.

Triactoma sp.

Figures 6.3, 6.4

Material.—Specimens P1-12-4, P1-12-8 are from the sample P1-012.

Description.—Cortical shell large, spherical, composed of very massive hexagonal or pentagonal pore frames; about twelve pore frames visible on test surface along axis of each spine. Three spines broken, two of them closely spaced, the third one widely separated from the other two.

Dimension (in μm).—Diameter of cortical shell: 228.

Remarks.—The specimens preserved with broken spines, two of them closely spaced, cortical shell large, spherical with the hexagonal and pentagonal pore frames.

Range and occurrence.—Upper Jurassic. Nadanhada.

Order Nassellaria Ehrenberg, 1875

Genus *Arcanica* Takemura, 1986

Type species.—*Arcanicapsa sphaerica* Takemura, 1986.

Arcanicapsa funatoensis (Aita, 1987)

Figures 9.16, 9.17

Sethocapsa funataoensis Aita, 1987, p. 73, pl. 2, figs. 6a–7b; pl. 9, figs. 14, 15;
Baumgartner *et al.*, 1995, p. 494, pl. 3070, figs. 1–4, 5.

Zhamoidellum funatoensis (Aita). Hull, 1997, p. 132, pl. 38, figs. 13, 15.

Arcanicapsa funatoensis (Aita, 1987). Nishihara and Yao, 2005, fig. 5.1; O’Dogherty
et al., 2017, p. 45, figs. 10.5–10.16.

Material.—Specimens P1-8-49, 59 are from the sample P1-08.

Dimensions (in μm).—Height/maximal width: 174/143 (P1-8-49), 153/158 (P1-8-59).

Remarks.—The Chinese specimens are slightly different from the Japanese type specimen by having roughened thorax with less number of pores.

Range and occurrence.—UAZones 3–11, early–middle Bajocian to late Kimmeridgian–early Tithonian (Baumgartner *et al.*, 1995). California, Austria, Japan, Nandanhada.

Genus *Archaeodictyomitra* Pessagno, 1976, emend. Pessagno, 1977b, Yang, 1993

Type species.—*Archaeodictyomitra squinaboli* Pessagno, 1976.

Archaeodictyomitra rigida Pessagno, 1977a

Figures 6.5, 6.6, 8.12, 8.13

Archaeodictyomitra rigida Pessagno, 1977a, p. 81, pl. 7, figs. 10, 11; Wu, 1993, p. 123, pl. 2, fig. 18; Hull, 1997, p. 79, pl. 32, fig. 6.

Material.—Specimens P1-8-38, P1-8-39 are from the sample P1-08; specimens P1-12-26, P1-12-43 are from the sample P1-012

Dimensions (in μm).—Height/Width: 152/84 (P1-8-38), 160/93 (P1-8-39), 182/99 (P1-12-26), 134/71 (P1-12-43).

Remarks.—The Chinese specimens have lesser number of chambers than the holotype of the species, but the other characters agree well with the type specimen.

Range and occurrence.—Middle Oxfordian–lower Kimmeridgian to Berriasian or younger. California; Nadanhada, Tibet, China.

Archaeodictyomitra sp.

Figure 8.14

Material.—Specimen P1-08-20 is from the sample P1-08.

Dimensions (in μm).—Length/Width: 167/101 (P1-08-20)

Remarks.—The specimens have the similar outline, and the same number of widely spaced longitudinal costae with one of the two figured specimens of

Archaeodictyomitra pseudomulticostata of Dumitrica *et al.* (1997, pl. 7, fig. 5), but differs in that the latter species has a inflated final chamber.

Range and occurrence.—Upper Jurassic. Nadanhada.

Genus *Caneta* Pessagno, Blome and Hull in Pessagno *et al.*, 1993

Type species.—*Parvicingula hsui* Pessagno, 1977a

Caneta? sp. 1

Figure 5.5

Material.—Specimen P1-09-22 is broken and from the sample P1-09.

Dimension (in μm).—Height/Maximum width: 217/125.

Remarks.—The Chinese specimen is preserved with broken cephalis, it has small and broader test than the type species *Caneta hsui*.

Range and occurrence.—Middle Jurassic. Nadanhada.

Caneta sp. 2

Figure 9.1

Material.—Specimen P108-1 is from the sample P1-08.

Dimension (in μm).—Height/Width (maximum): 308/131.

Remarks.—The Chinese specimen is similar to *Svinitzium pseudopuga* Dumitrica,

1997, but differs by having slender test outline, and stronger H-linked costae in the circumferential ridges.

Range and occurrence.—Upper Jurassic. Nadanhada.

Genus *Cinguloturris* Dumitrica in Dumitrica and Mello, 1982

Type species.—*Cinguloturris carpatica* Dumitrica in Dumitrica and Mello, 1982.

Cinguloturris fusiforma Hori, 1999

Figure 8.19

Cinguloturris fusiforma Hori, 1999, p. 93, fig. 9(3–6); fig. 11(6).

Material.—Specimen P1-08-4 is from the sample P1-08.

Dimensions (in μm).—Height/Maximum width: 302/163.

Remarks.—The species differs from *Cinguloturris carpatica* by having spindle shape outline.

Range and occurrence.—Tithonian (Upper Jurassic). Russia, Japan, Nadanhada.

Cinguloturris carpatica Dumitrica in Dumitrica and Mello, 1982

Figures 6.7, 8.20, 8.21

Cinguloturris carpatica Dumitrica. Dumitrica and Mello, 1982, p. 23, pl. 4, figs.,

7–14; Matsuoka and Yao, 1986, pl. 2, fig. 14; Matsuoka, 1986a, pl. 2, fig. 16; pl. 3, fig. 11a, b; Baumgartner *et al.*, 1995, p. 142, pl. 3193, figs. 1–6; Yang and Matsuoka, 1997, pl. 3, fig. 9; Hori, 1999, p. 91, fig. 9(1); Matsuoka *et al.*, 2002, fig. 5(16).

Material.—Specimens P1-8-4, P1-8-9, P1-8-43, P1-8-46 are from the sample P1-08; specimen P1-12-11 is from the sample P1-012.

Dimensions (in μm).—Height/Maximum width: 189/108 (P1-8-4), 252/102 (P1-8-9)

Remarks.—Diameter of segment increasing rapidly up to the third segment, and slower in the following ones, so that the postabdominal chambers become cylindrical. The spongy network between adjacent segments is well developed beginning with the third or fourth constriction

Range and occurrence.—The species is ranged within UA Zones 7–11, upper Bathonian–lower Callovian to upper Kimmeridgian–lower Tithonian (Baumgartner *et al.*, 1995); Callovian to Tithonian (Hori, 1999). Caucasus, Japan, Nadanhada, northern Apennines, Russian Far East, Tibet, western Carpathians, western Pacific Ocean (ODP Leg 129, Site 801), Yugoslavia.

Genus *Crococapsa*

Type species.—*Tetracapsa pilula* Rüst, 1885

Crococapsa hexagona (Hori, 1999)

Figures 9.12–9.15

Sethocapsa hexagona Hori, 1999, p. 74, fig. 6(12–16); fig. 11(1a, b).

Material.—Specimens P1-8-10, 15, 19, 60, and P108-27 are from the sample P1-08.

Description.—Test consists of four segments. Cephalis subspherical, imperforate, without apical horn, and slightly encased in the thorax. Thorax and abdomen trapezoidal and perforate. First three segments are almost the same height and gradually increase in width. Strictures are weakly developed. The final segment is very large and spherical, and possesses strong hexagonal pore frames with rounded pores.

Dimensions (in μm).—Height/Maximal Width: 162/118 (P1-8-10), 149/115 (P1-8-15), 143/126 (P1-8-19), 142/124 (P1-8-60), ?/177 (P108-27).

Remarks.—The present specimens differ from the Japanese specimens of *T. hexagona* Hori by having smaller test, and by possessing a spherical final chamber, but with respect to the strong hexagonal pore frames, they can be assigned to the Japanese species. The present specimens are similar to *Hiscocapsa asseni* (Tan) (O’Dogherty, 1994) but differs from the latter by possessing larger and less number of pore frame in the shell of the final chamber, and by lacking a terminal aperture.

Range and occurrence.—Upper Jurassic. Japan, Nandanhada.

Genus *Eucyrtidiellum* Baumgartner, 1984

Type species.—*Eucyrtidium* (?) *unumaensis* Yao, 1979.

Eucyrtidiellum ptyctum (Riedel and Sanfilippo, 1974)

Figure 8.10

Eucyrtidium ptyctum Riedel and Sanfilippo, 1974, p. 778, pl. 5, fig. 7; pl. 12, fig. 14, not fig. 15; Matsuoka and Yao, 1985, pl. 2, fig. 8.

Eucyrtidiellum ptyctum (Riedel and Sanfilippo). Matsuoka, 1986a, pl. 2, fig. 10; Baumgartner *et al.*, 1995, p. 214, pl. 3017, figs. 1–8.

Material.—Specimen 108x2-24 is from the sample P1-08.

Dimensions (in μm).—Height/Maximum width: 95/87.

Remarks.—The present species differs from other species of the genus by having a thorax with numerous small nodes and wholly costated abdomen.

Range and occurrence.—UAZones 5–11, latest Bajocian–early Bathonian to late Kimmeridgian–early Tithonian (Baumgartner *et al.*, 1995). Callovian to Tithonian (Hori, 1999). Argolis Peninsula, Blake-Bahama Basin (DSDP Site 534), California Coast Ranges, western Pacific Ocean (ODP Leg 129, Site 801), Yugoslavia.

Eucyrtidiellum unumaense (Yao, 1979)

Figure 8.11

Eucyrtidium ptyctum Riedel and Sanfilippo, 1974, p. 778, pl. 12, fig. 15 only.

Eucyrtidium (?) *unumaensis* Yao, 1979, p. 39, pl. 9, figs. 1–11.

Eucyrtidium (?) *unumaense* Yao. Matsuoka, 1982, pl. 1, fig. 15; Matsuoka, 1985, pl. 1, fig. 9; Kojima, 1989, pl. 2, fig. 5a, b; Matsuoka, 1992, pl. 1, fig. 8; pl. 2, fig. 7.

Eucyrtidiellum unumaense unumaense (Yao). Baumgartner *et al.*, 1995, p. 222, pl. 3012, figs. 1, 2.

Material.—Specimen P1-8-64 is from the sample P1-08.

Dimension (in μm).—Height/Maximum width: 168/101.

Remarks.—The Chinese specimen is poorly preserved, and with relatively long thorax, but these may be intraspecific variation.

Range and occurrence.—UAZones 3–8, early-middle Bajocian to middle Callovian–early Oxfordian (Baumgartner *et al.*, 1995). Far East Russia, Japan, Yugoslavia, Nadanhada.

Genus *Gongylothorax* Foreman, 1968, emend. Dumitrica, 1970

Type species.—*Dicolocapsa verbeeki* Tan Sin Hok, 1927.

Gongylothorax favosus Dumitrica, 1970

Figure 8.9

Gongylothorax favosus Dumitrica, 1970, p. 56, pl. 1, fig. 1a–c, 2; Baumgartner *et al.*, 1995, p. 230, pl. 6131, figs. 1–6, 7; Hori, 1999, p. 74, fig. 6(10).

Material.—Specimen P1-08-11 is from the sample P1-08.

Dimensions (in μm).—Width of thorax: 153.

Remarks.—The Chinese specimen with a spherical thorax, whose surface are ornamented with regularly arranged hexagonal pore frames, bearing a small pore in the center. All these agree well with *Gongylothorax favosus*.

Range and occurrence.—UAZones 8–10, mid Callovian–early Oxfordian to late Oxfordian–early Kimmeridgian (Baumgartner *et al.*, 1995). Bathonian to Tithonian (Hori, 1999). California, Eastern Alps, Japan, Nadanhada, Romania, western Pacific Ocean (ODP Leg 129, Site 801), Yugoslavia.

Genus *Loopus* Yang, 1993

Type species.—*Pseudodictyomitra primitiva* Matsuoka and Yao, 1985.

Loopus sp. 1

Figure 8.15

Material.—Specimen P1-08-2 is from the sample P1-08.

Dimensions (in μm).—Length/Width: 265/103.

Remarks.—The present specimen is attributed to *Loopus* because most segments bear longitudinal and discontinuous costae.

Range and occurrence.—Upper Jurassic. Nadanhada.

Loopus sp. 2

Figures 9.3–9.5

Material.—Specimens P1-08-30, 108x2-11, 108x2-12 are from sample P1-08.

Dimension (in μm).—Height/Maximal width: 181/97 (P1-08-30), 164/99 (108x2-11), 209/108 (108x2-12).

Remarks.—The Chinese specimens are similar to *Loopus nudus* (Schaaf, 1981) (Dumitrica *et al.*, 1997, pl. 5, figs. 6, 23) in having the same test outline, the same number of chambers, but the former differs from the latter by having weakly developed strictures between chambers, and by having more distinct fine costae on chamber surface.

Range and occurrence.—Upper Jurassic. Nadanhada.

Loopus? sp. 3

Figures 9.6, 9.7

Material.—Specimens P1-08-6, 25 are from the sample P1-08.

Dimensions (in μm).—Height/Maximal width: 169/107 (P1-08-6), 163/94 (P1-08-25).

Remarks.—The Chinese specimens possess one row of large primary pores in strictures between segments, but the postcephalic chambers possess three rows of primary and relict pores instead of fine costae. They differ from the typical taxa of

Loopus, thus they are tentatively assigned to the genus *Loopus*. The Chinese specimens are closely similar to *Pseudodictiomitra?* sp. D from the Upper Jurassic to Lower Cretaceous Torinosu Group and Yura Formation (Matsuoka and Yao, 1985, pl. 2, figs. 6, 7), and from the Upper Jurassic of Togano Group (Matsuoka, 1986a, pl. 2, fig. 11) of Japan, but the former differs from the latter in having three rows of primary and relict pores in each postcephalic segment instead of two rows in the latter taxon.

Range and occurrence.—Upper Jurassic. Nadanhada.

Genus *Olanda* Hull, 1997

Type species.—*Olanda olorina* Hull, 1997.

Olanda? sp.

Figure 9.2

Material.—One broken specimen 108x2-25 is from the sample P1-08.

Dimension (in μm).—Height/Maximal width: 229/127.

Remarks.—The present broken specimen is composed of an apical slender conical part and a large final postabdominal bulbous chamber. But the apical conical part has not longitudinal costae, which is well developed in the type species *Olanda olorina*.

The Chinese specimen is similar to *Sethocapsa dorysphaeroides* Neviani (Baumgartner et al., 1995, p. 494, pl. 5544), but differs from the latter by having constricted aperture.

Range and occurrence.—Upper Jurassic. Nadeshima.

Genus *Protunuma* Ichikawa and Yao, 1976

Type species.—*Protunuma fusiformis* Ichikawa and Yao, 1976.

Protunuma japonicus Matsuoka and Yao, 1985

Figures 8.17, 8.18

Protunuma fusiformis Ichikawa and Yao. Mizutani, 1981, p. 181, pl. 63, figs. 1, 8; pl. 64, fig. 3.

Protunuma japonicus Matsuoka and Yao, 1985, p. 130, pl. 1, figs. 11–15; pl. 3, figs. 6–9; Matsuoka, 1986a; pl. 2, fig. 7; Baumgartner *et al.*, 1995, p. 434, pl. 3292, figs. 1–8; Hull, 1997, p. 156, pl. 43, figs. 8, 14, 15; Zügel, 1997, p. 204, fig. 5(8); Hori, 1999, p. 85, fig. 7(20); Kiessling, 1999, p. 28, pl. 13, fig. 19; Matsuoka *et al.*, 2005, figs. 4(11), 5(11).

Material.—Specimens P1-8-1, P1-8-25 are from the sample P1-08.

Dimension (in μm).—Height/Width: 174/115 (P1-8-1), 183/120 (P1-8-25)

Remarks.—This species is similar to *Protunuma fusiformis* Ichikawa and Yao and *P. turbo* Matsuoka in outer shape, but differs in that the latter two forms have externally recognizable lumbar stricture, because the abdomen expands more strongly than thorax in the latter two species.

Range and occurrence.—UAZones 7–12, late Bathonian–early Callovian to early–early late Tithonian (Baumgartner et al., 1995); Oxfordian–Tithonian (Hori, 1999). Blake-Bahama Basin (DSDP Site 534), California Coast Ranges, Eastern Alps, Eastern Atlantic Ocean (DSDP Leg 41, Site 367), Japan, Nadanhada, Northern Apennines, Northern Calcareous Alps, Russian Far East, Solnhofen (Southern Germany), Southern Alps, Southern Tibet, Western Carpathians, western Pacific Ocean (ODP Leg 129, Site 801), Yugoslavia.

Genus *Pseudoxitus* Wu and Pessagno in Wu, 1993; emend. Dumitrica in Dumitrica et al., 1997

Type species.—*Pseudoxitus inflatus* Wu, 1993.

Pseudoxitus? sp.

Figure 6.10

Material.—Specimen P1-12-19 is from the sample P1-012.

Dimension (in μm).—Height/Maximal width: 207/160.

Remarks.—The present form is tentatively attributed to *Pseudoxitus* because of lacking well developed tubercles. The outer layer structure mainly consists of irregular bars. The specimen differs the known taxa of the genus by the cylindrically inflated final chamber.

Range and occurrence.—Middle Jurassic. Nadanhada.

Genus *Ristola* Pessagno and Whalen, 1982, emend. Baumgartner, 1984

Type species.—*Parvicingula* (?) *procera* Pessagno, 1977a.

Ristola sp.

Figure 7.4

Material.—Specimen P1-13-11 is from the sample P1-013.

Description.—Test preserved with the apical part. Cephalis with a massive horn. Apical part covered by a outer layer of broad bars, the inner layer of test invisible.

Dimensions (in μm).—Height/Maximal width: 180/155.

Remarks.—Only one broken specimen is encountered, because of the conical, apical portion is covered by a outer layer it indicates that the specimen could be assigned to

Ristola.

Range and occurrence.—Upper Jurassic, Nadanhada.

Genus *Solenotryma* Foreman, 1968

Type species.—*Solenotryma dacryodes* Foreman, 1968

Solenotryma ichikawai Matsuoka and Yao, 1985

Figure 6.9

Solenotryma? ichikawai Matsuoka and Yao, 1985, p. 133, pl. 3, fig. 21.

Solenotryma ichikawai Matsuoka and Yao. Baumgartner *et al.*, 1995, p. 508, pl. 4037, figs. 1–5; Hori, 1999, p. 87, fig. 8.12; Matsuoka *et al.*, 2005, fig. 5(12).

Material.—Specimen P1-12-41 is from the sample P1-012.

Dimensions (in μm).—Height/Maximal width: 193/112.

Remarks.—The Chinese specimen differs from the species *Solenotryma palmerae* Pessagno in that the latter possesses more chambers and has a slenderer test outline. The present specimen differs from *S. perampla* (Rüst) in that the latter being much larger forms with more broader conical test outline.

Range and occurrence.—UAZones 7–21, late Bathonian–early Callovian to early Barremian (Baumgartner *et al.*, 1995); Torinosu Group, southwestern Japan; Nandanhada.

Genus *Stichocapsa* Haeckel, 1881

Type species.—*Stichocapsa jaspidea* Rüst, 1885.

“*Stichocapsa*” *robusta* Matsuoka, 1984

Figures 5.7, 5.8

Stichocapsa robusta Matsuoka, 1984, p. 146, pl. 1, figs. 6–13; pl. 2, figs. 7–12;

Matsuoka, 1986b, pl. 1, fig. 12; Matsuoka, 1988, pl. 1, fig. 8; Matsuoka, 1992, pl.

5, fig. 3; Baumgartner *et al.*, 1995, p. 524, pl. 3298, figs. 1–7; Matsuoka *et al.*, 2002, fig. 5(4).

Material.—Specimens P1-09-24 and 9-5 are from the sample P1-09.

Dimensions (in μm).—Height/Maximal width: 165/126 (P1-9-5); 232/181 (P1-09-24).

Remarks.—The lumbar sutural pore not visible. The species is similar to “*Stichocapsa*” *convexa* Yao, 1979, but differs from the latter in its thick wall, internally tapering pores and wider apical conical part.

Range and occurrence.—Middle to lower Upper Jurassic. Nadanhada and Tibet, China; Japan; southern California Coast Ranges.

“*Stichocapsa*” *convexa* Yao, 1979

Figures 5.9, 5.10

Stichocapsa convexa Yao, 1979, p. 35, pl. 5, figs. 14–16; pl. 6, figs. 1–7; Matsuoka, 1985, pl. 1, fig. 8; Baumgartner *et al.*, 1995, p. 518, pl. 3055, figs. 1–4.

Material.— Specimen P1-09-10 is from the sample P1-09.

Dimensions (in μm).—Height/Maximal width: 225/170.

Remarks.—Shell of four segments, cephalis poreless, thorax and abdomen truncated-cornical. The fourth segment truncate-spherical with small aperture.

Range and occurrence.—Middle and Upper Jurassic. Nadanhada; Japan.

“*Stichocapsa*” sp.

Figure 5.11

Material.—Specimen P1-9-8 is from the sample P1-09.

Dimensions (in μm).—Height/Maximal width: 131/110.

Remarks.—The Chinese specimen differs from the species *Stichocapsa praepulchella* Hori, 1999 by having fewer rows of transverse pores, less number of segments and smaller test.

Range and occurrence.—Upper Jurassic. Nadanhada.

Genus *Striatojaponocapsa* Kozur, 1984, emend. Hull, 1997

Type species.—*Tricolocapsa plicarum* Yao, 1979.

Striatojaponocapsa synconexa O'Dogherty, Gorican and Dumitrica in O'Dogherty *et al.*, 2005

Figures 5.12, 5.13

Tricolocapsa plicarum Yao. Matsuoka, 1983, p. 20, pl. 3, fig. 2.

Tricolocapsa plicarum ssp. A. Baumgartner *et al.*, 1995, p. 598, pl. 4052, figs. 1–5.

Striatojaponocapsa synconexa O'Dogherty, Gorican and Dumitrica in O'Dogherty *et al.*, 2005, p. 447, pl. 10, figs. 9–17.

Material.—Specimen P1-9-28 is from the sample P1-09.

Dimensions (in μm).—Height/Maximal width: 140/132.

Remarks.—Outer surface of shell ornamented with continuous longitudinal plicae, without transverse ridges between adjacent plicae.

Range and occurrence.—UAZones 4–5, late Bajocian–early Bathonian (Baumgartner *et al.*, 1995), Japan, Greece, eastern Alps, western Carpathian, Nadezhda; Middle to Upper Jurassic of the United States.

Genus *Tethysetta* Dumitrica in Dumitrica *et al.*, 1997

Type species.—*Tethysetta pygmaea* Dumitrica in Dumitrica *et al.*, 1997.

Tethysetta dhimenaensis (Baumgartner, 1984)

Figures 6.8, 8.22–8.25

Parvicingula dhimenaensis Baumgartner, 1984, p. 778, pl. 7, figs. 2, 3; Matsuoka, 1986a, pl. 2, fig. 12; Dumitrica *et al.*, 1997, p. 49, pl. 11, figs.

Parvicingula dhimenaensis dhimenaensis Baumgartner. Baumgartner *et al.*, 1995, p. 406, pl. 4072, figs. 1–3; Chiari *et al.*, 1997, pl. 3, fig. 3; Zügel, 1997, p. 206, fig. 5(13); Hori, 1999, p. 96, fig. 9(16).

Tethysetta dhimenaensis (Baumgartner). Nishida, 2011, p. 108, fig. 6.19.

Material.—Specimens 108x2–22, 28, P1-8-24, 27, 37, 58, 68 are from the sample P1-08; specimens P1-12-9, 15, 24, 27 are from the sample P1-012.

Dimensions (in μm).—Height/Maximum width: 240/113 (108x2-28), 163/89 (P1-8-27), 161/95 (P1-8-37), 212/91 (P1-8-42), 195/96 (P1-8-58), 228/113 (P1-8-68), 225/105 (P1-12-9), 296/131 (P1-12-15), 231/110 (P1-12-24).

Remarks.—Baumgartner (1984) described the species *Parvicingula dhimenaensis* from Greece, which including at least two morphotypes. According to the character of the proximal portion of tests, Baumgartner *et al.* (1995) separated these two morphotypes into two subspecies, i.e., *P. dhimenaensis dhimenaensis* and *P. dhimenaensis* ssp. A. The former has shorter cephalo-thorax portion, while the latter with long and externally smooth proximal portion which is composed of cephalis, thorax, abdomen and possibly the first postabdominal chamber. As noted by Baumgartner (1984) the species *P. dhimenaensis* differs from other *Parvicingula* in having circumferential ridges with regularly spaced nodes and diagonal bars connecting between nodes. Thus Dumitrica *et al.* (1997) attributed this species to *Tethysetta*. The Chinese specimens agree very well with the subspecies *Tethysetta dhimenaensis*.

Range and Occurrence.—UAZones 3–11 (early-mid Bajocian to early Tithonian) (Baumgartner *et al.*, 1995), or Callovian to Tithonian (Hori, 1999), up to Berriasian (Kiessling, 1995; Dumitrica *et al.*, 1997). Nadanhada, east-central Mexico, Japan, Northern Apennines, Solnhofen (southern Germany), Southern Alps, Western Carpathians, Western Himalaya, Yugoslavia.

Genus *Theocapsomella* O'Dogherty, Gorican and Dumitrica in O'Dogherty *et al.*,
2005

Type species.—*Theocapsomma cordis* Kocher, 1981

Theocapsomella himedaruma (Aita, 1987)

Figure 9.18

Stichocapsa himedaruma Aita, 1987, p. 74, pl. 3, figs. 1a–3; pl. 10, figs. 1, 2;
O'Dogherty *et al.*, 2005, p. 452.

Material.—Specimen P1-8-54 is from the sample P1-08.

Dimensions (in μm).— Height/Maximal width: 102/91.

Remarks.—The Chinese specimen possesses the similar test outline as the Japanese species *Theocapsomella himedaruma*, but differs by having a relatively smaller abdomen and a larger final segment. This difference may be intraspecific variation.

Range and occurrence.—Late Middle–early Late Jurassic. Japan, Nandanhada.

Genus *Transhsuum* Takemura, 1986

Type species.—*Transhsuum medium* Takemura, 1986

Transhsuum? sp.

Figure 5.2

Material.—Specimen P1-9-3 is from the sample P1-09.

Dimensions (in μm).—Length/Width: 264/108.

Remarks.—The present specimen is questionably attributed to *Transhsuum* because it is multi-segmented nassellaria, but discontinuous longitudinal costae are poorly preserved.

Range and occurrence.—Middle Jurassic. Nanhada.

Genus *Tricolocapsa* Haeckel, 1881

Type species.—*Tricolocapsa theophrasti* Haeckel, 1887

“*Tricolocapsa*” sp.

Figure 5.14

Material.—Specimen P1-9-20 is from the sample P1-09.

Dimensions (in μm).—Height/Maximal width: 129/120.

Remarks.—Test with three segments. Shell surface ornamentation not preserved.

Range and occurrence.—Middle Jurassic. Nanhada.

“*Tricolocapsa*” *tetragona* Matsuoka, 1983

Figures 5.15–5.16

Tricolocapsa tetragona Matsuoka, 1983, p. 22, pl. 3, figs. 8–12; pl. 8, figs. 4–10;
Matsuoka and Yao, 1986, pl. 1, fig. 18; Baumgartner *et al.*, 1995, p. 600, pl. 4054,
figs. 1–3.

Material.—Specimens P1-9-6 is from the sample P1-09.

Description.—Test with three segments. Cephalis small, rounded, poreless. Thorax cylindrical with small pores. Abdomen spherical with longitudinal plicae and transverse ridges connecting adjacent two longitudinal plicae.

Dimensions (in μm).—Height/Maximal width: 175/170.

Remarks.—The present specimen is poorly preserved with several longitudinal plicae and transverse ridges between plicae on the abdomen. It is possible to be assigned to

Tricolocapsa tetragona.

Range and occurrence.—UAZone 5 (latest Bajocian–early Bathonian) (Baumgartner *et al.*, 1995). Japan, Nandahada.

Genus *Unuma* Ichikawa and Yao, 1976

Type species.—*Unuma typicus* Ichikawa and Yao, 1976.

Unuma gordus Hull, 1997

Figures 5.3, 5.4, 8.16

Unuma sp. A. Baumgartner *et al.*, 1995, p. 624, pl. 3309, figs.1–4.

Unuma gorda Hull, 1997, p. 172, pl. 43, figs. 9, 11, 12; Chiari *et al.*, 2003, pl. 2, fig. 19; Chiari *et al.*, 2004, pl. 3, fig. 15.

Unuma gordus Hull. Suzuki and Gawlick, 2009, p. 177, fig. 6(2A, B); Djerić *et al.*, 2012, p. 362, pl. 1, fig. 24.

Material.—Specimen P1-8-69 is from the sample P1-08; Specimens P1-9-4, P1-9-7 are from sample P1-09..

Dimensions (in μm).—Height/Width: 191/139 (P1-8-69), 120/92(P1-9-4), 137/107(P1-9-7).

Remarks.—The genus *Unuma* was established as masculine gender (Ichikawa and Yao, 1976), so that the species name was changed to *gordus* by Suzuki and Gawlick (2009). Test broadly spindle-shaped, with four segments. The present specimens are closely related to *Unuma* sp. A (Baumgartner *et al.*, 1995, p. 624, pl. 3309, figs. 1–4), which is considered a synonym of *Unuma gordus*.

Range and occurrence.—UAZones 4–6, late Bajocian–mid Bathonian (Baumgartner *et al.*, 1995), United States, Japan, southern Alps, Nanhada.

Genus *Willriedellum* Dumitrica, 1970.

Type species. —*Willriedellum crystallinum* Dumitrica, 1970

Willriedellum marcucciae (Cortese, 1993)

Figures 9.10, 9.11

Williriedellum sp. A gr. Matsuoka, 1983, p. 23, pl. 4, figs. 1–3; pl. 8, figs. 11–15.

Williriedellum marcuccii Cortese, 1993, p. 180, pl. 7, figs. 6, 7.

Williriedellum marcucciae (Cortese, 1993). Suzuki and Gawlick, 2009, p. 179, fig. 5(25), fig. 6(49A, B).

Material.—Specimen P1-8-40 is from the sample P1-08.

Dimensions (in μm).—Height/Maximal width: 169/131.

Remarks.—The species name has been revised from *marcuccii* to *marcucciae* because the species was named after Prof. Marta Marcucci (Suzuki and Gawlick, 2009).

Range and occurrence.—UAZones 4–8, late Bajocian–early Oxfordian (Baumgartner et al., 1995). Austria, Japan, Nanhada.

Genus *Zhamoidellum* Dumitrica, 1970

Type species.—*Zhamoidellum ventricosum* Dumitrica, 1970.

Zhamoidellum ovum Dumitrica, 1970

Figures 6.11, 9.8, 9.9

Zhamoidellum ovum Dumitrica, 1970, p. 79, pl. 9, figs. 52a, b, 53, 54; Baumgartner et al., 1995, p. 656, pl. 4079, figs. 2, 4, 6; O’Dogherty et al., 2017, p. 50,

figs.10.31–10.40.

Material.—Specimens P1-8-18, 32, 50, 55 are from the sample P1-08; specimen P1-12-5, 30 are from the sample P1-012.

Dimensions (in μm).—Height/Maximal width: 145/142 (P1-8-18), 128/116 (P1-8-32), 128/97 (P1-8-47), 119/108 (P1-8-50), 109/93 (P1-8-55), ?/173 (P1-12-5), 194/146 (P1-12-30).

Remarks.—Baumgartner *et al.* (1995) figured six specimens for the species *Zhamoidellum ovum*, their specimen (pl. 4079, fig. 1) is closely related to *Z. mikamense* Aita, 1987 because of the well developed lumbar stricture; the specimens (pl. 4079, figs. 3 and 5) show dimple-shaped pits in the lumbar stricture, within the pits there are smaller pore frames than the surrounding ones, these pits look like sutural pores, but Dumitrica has not mentioned this structure in the original description of the species, thus these two specimens should be assigned to the genus *Complexapora* Kiessling in Kiessling and Zeiss, 1992.

The Chinese specimens are closely related to *Z. ovum*, not only in test outline, but also possess the same porous structure in the thick-walled thorax and abdomen.

Range and occurrence.—UAZones 9–11, mid–late Oxfordian to early Tithonian (Baumgartner *et al.*, 1995). Romania, Japan, Nadanhada.

***Zhamoidellum* sp.**

Figure 9.11

Material.—Specimens P1-8-44 is from the sample P1-08.

Dimensions (in μm).—Height/Maximal width: 164/133.

Remarks.—The thorax porous with the lower part constricted and depressed into the abdominal cavity, showing a clear lumbal stricture.

Range and occurrence.—Upper Jurassic, Nadanhada.

Conclusion

Four black claystone samples (P1-08, 09, 012, 013) from the Upper Jurassic Dalingqiao Formation of the Nadanhada terrane contain well-preserved radiolarian faunas. The sample P1-09 yields a *Striatojaponocapsa synconexa*-“*Tricolocapsa*” *tetragona* assemblage, which indicates a Middle Jurassic Bathonian age. The other three samples contain a Late Jurassic *Cinguloturris carpatica* assemblage, which may be of a late Oxfordian–early Tithonian age. The discovery of new Late Jurassic radiolarians indicates that the Raohe Complex was accreted during the Late Jurassic paleo-Pacific subduction. Then, the succedent late Tithonian–early Valanginian *Buchia* fauna-bearing Dong’anzen Formation may be the earliest cover beds overlying the Raohe Complex. The development of the Lower Cretaceous *Aucellina* bivalve fauna-bearing Dajiashan Group indicates a paleogeographic differentiation between the Nadanhada terrane and the Tamba-Mino-Ashio terrane.

Acknowledgements

This study was supported by the Strategic Priority Research Program of Chinese Academy of Sciences (XDB26000000), by the National Natural Science Foundation of China (41572006, 41688103, 41781240013, 91514302, 41172010). Great thanks go to Prof. Peiji Chen of Nanjing Institute of Geology and Palaeontology, Chinese Academy of Sciences for his supervision and planning for this research. We are grateful for the guide of Prof. Baoyu Jiang (Nanjing University, China) to the field sections of the Dajiashan Group. Many thanks go to Prof. Keisuke Ishida (University of Tokushima, Japan) and an anonymous reviewer for their constructive review. It is also a contribution to UNESCO-IUGS IGCP Project 679.

References

- Aita, Y., 1987: Middle Jurassic to Lower Cretaceous radiolarian biostratigraphy of Shikoku with reference to selected sections in Lombardy Basin and Sicily. *Science Reports of the Tohoku University, 2nd Series (Geology)*, vol. 58, p. 1–91.
- Baumgartner, P. O., 1980: Late Jurassic Hagiastriidae and Patulibracchiidae (Radiolaria) from the Argolis Peninsula (Peloponnesus, Greece). *Micropaleontology*, vol. 26, p. 274–322.
- Baumgartner, P. O., 1984: A Middle Jurassic–Early Cretaceous low latitude radiolarian zonation based on unitary associations and age of Tethyan radiolarites. *Eclogae Geologicae Helvetiae*, vol. 77, p. 729–841.

Baumgartner, P. O., Dogherty, L. O', Gorican, Š., Dumitrica-Jud, R., Dumitrica, P., Pillecuit, A., Urquhart, E., Matsuoka, A., Danelian, T., Bartolini, A., Carter, E. S., De Wever, P., Kito, N., Marcucci, M. and Steiger, T., 1995: Radiolarian catalogue and systematics of Middle Jurassic to Early Cretaceous Tethyan genera and species. In, Baumgartner, P. O., Dogherty, L. O', Gorican, Š., Urquhart, E., Pillecuit, A., De Wever, P. eds., *Middle Jurassic to Lower Cretaceous Radiolaria of the Tethys: occurrences, systematics, biochronology. Mémoires de Géologie (Lausanne)*, no. 23, p. 37–688.

Bureau of Geology and Mineral Resources of Heilongjiang Province, 1993: Regional Geology of Heilongjiang Province. *Geological Memoirs, 1st series*, vol. 33, p. 1–734. (in Chinese with English summary)

Buryi, G.I., 1996: Triassic conodonts from the cherts of Nadanhada range, northeast China. *Acta Micropalaeontologica Sinica*, vol. 13, p. 207–214.

Campbell, A. S., 1954: Radiolaria. In, Moore, R. C. ed., *Treatise on Invertebrate Paleontology. Part D, Protista 3*, p. 11–195. Geological Society of America and University of Kansas Press, Lawrence.

Chiari, M., Bortolotti, V., Marcucci, M., Photiades, A. and Principi, G., 2003: The Middle Jurassic siliceous sedimentary cover at the top of the Vourinos ophiolite (Greece). *Ofioliti*, vol. 28, p. 95–103.

Chiari, M., Corstese, Marcucci, M. and Nozzoli, N., 1997. Radiolarian biostratigraphy in the sedimentary cover of the ophiolites of south-western Tuscany, Central Italy. *Eclogae Geologicae Helvetiae*, 90, p. 55-77.

- Chiari, M., Marcucci, M. and Prela, M., 2004: Radiolarian assemblages from the Jurassic cherts of Albania: new data. *Ofioliti*, vol. 29, p. 95–105.
- Cortese, G., 1993: Radiolarian biostratigraphy of the Tuscan Cherts (Tuscan Succession) from Val di Lima, Tuscany, Northern Apennines. *Paleopelagos*, vol. 3, p. 165–185.
- De Wever, P., Dumitrica, P., Caulet, J.P., Nigrini, C., Caridroit, M., 2001. Radiolarians in the Sedimentary Record. Gordon and Breach Science Publishers, 525 pp.
- Djerić, N., Schmid, S. M. and Gerzina, N., 2012: Middle Jurassic radiolarian assemblages from the sedimentary cover of the Adriatic margin (Zlatar Mountain, SW Serbia). *Bulletin de la Société Géologique de France*, vol. 183, p. 359–368.
- Dumitrica, P., 1970: Cryptocephalic and cryptothoracic Nasselaria in some Mesozoic deposits of Romania. *Revue Roumaine de Géologie, Géophysique et Géographie (série Géologie)*, vol. 14, p. 45–124.
- Dumitrica, P. and Mello, J., 1982: On the age of the Meliata Group and the Silica Nappe radiolarites (localities Drzkovce and Bohunovo, Slovak Karst, CSSR). *Geologické Práce*, vol. 77, 17–28.
- Dumitrica, P., Immenhauser, A. and Dumitrica-Jud, R., 1997: Mesozoic radiolarian biostratigraphy from Masirah Ophiolite, Sultanate of Oman. Part I: Middle Triassic, Uppermost Jurassic and Lower Cretaceous Spumellarians and multisegmented Nassellarians. *Bulletin of the National Museum of Natural Science*, vol. 9, p. 1–106.
- Ehrenberg, C. G., 1875: Fortsetzung der mikrogeologischen Studien als

Gesamt-Übersicht der mikroskopischen Paläontologie gleichartig analysirter Gebirgsarten der Erde, mit specieller Rücksicht auf den Polycystinen-Mergel von Barbados. *Königliche Akademie der Wissenschaften zu Berlin, Abhandlungen, Jahre 1875*, p. 1–225.

Foreman, H. P., 1968: Upper Maestrichtian Radiolaria of California. *Special Papers in Palaeontology*, vol. 3, 1–82.

Foreman, H. P., 1973: Radiolaria from DSDP Leg 20. In, Heezen, B. C., MacGregor, I. D., Foreman, H. P. *et al. eds.*, *Initial Report of the Deep Sea Drilling Project*, vol. 20, p. 249–305, pl. 1–16.

Futakami, M., Matsukawa, M., Chen, P. J. and Chen, J. H., 1995: Barremian ammonites from the Longzhaogou Group in eastern Heilongjiang, northeast China. *Journal of the Geological Society of Japan*, vol. 101, p. 79–85.

Haeckel, E., 1881: Entwurf eines Radiolarien-Systems auf Grund von Studien der Challenger-Radiolarien. *Jenaische Zeitschrift für Naturwissenschaft*, 15 new serie, Band 8, p. 418–472.

Haeckel, E., 1887: Report on the Radiolaria collected by H.M.S. Challenger during the years 1873–1876. *Report on the Scientific Results of the Voyage of the H.M.S. Challenger, Zoology*, vol. 18, clxxxviii+1803.

Hori, N., 1999: Latest Jurassic radiolarians from the northeastern part of the Torinoko Bolck, Yamizo Mountains, central Japan. *Science Report of the Institute of Geoscience, University of Tsukuba, Section B, Geological Sciences*, vol. 20, p. 47–114.

- Hull, D. M., 1997: Upper Jurassic Tethyan and southern Boreal radiolarians from western North America. *Micropaleontology*, vol. 43, supplement 2, p. 1–202, pl. 1–51.
- Ichikawa, K. and Yao, A., 1976: Two new genera of Mesozoic cyrtoid radiolarians from Japan. In, Takayanagi, Y., Saito, T. eds., *Progress in Paleontology, Special Publication*, p. 110–117. Micropaleontology Press, the American Museum of Natural History, New York.
- Ishida, N., 2011: An exotic body of Middle Jurassic Torinosu-type limestone in the Hikawa Formation, southeastern Kanto Mountains, Japan. *Memoir of the Fukui Prefectural Dinosaur Museum*, vol. 10, p. 103–112.
- Kiessling, W., 1995: New radiolarians from the Earliest Cretaceous of the Sultanate of Oman (Wahrah Formation, Jebel Buwaydah). *Palaeontologisch Zeitschrift*, vol. 69, p. 321–342.
- Kiessling, W., 1999: Late Jurassic radiolarians from the Antarctic Peninsula. *Micropaleontology*, vol. 45, supplement 1, p. 1–96.
- Kiessling, W. and Zeiss, A., 1992: New palaeontological data from the Hochstegen Marble (Tauern window, eastern Alps). *Geologisch Paläontologische Mitteilungen Innsbruck*, vol. 18, p. 187–202.
- Kocher, R. N., 1981: Biochronostratigraphische Untersuchungen oberjurassischer radiolarien führender Gesteine, insbesondere der Südalpen. *Mitteilungen aus dem geologischen Institut der Eidgenössischen Technischen Hochschule und der Universität Zürich, Neue Folge*, Band 234, p. 1–184.
- Kojima, S., 1989: Mesozoic terrane accretion in northeast China, Sikhote-Alin and

- Japan regions. *Palaeogeography, Palaeoclimatology, Palaeoecology*, vol. 69, p. 213–232.
- Kojima, S. and Kametaka, M., 2000: Jurassic accretionary complexes in East Asia. *Memoirs of the Geological Society of Japan*, no. 52, p. 61–72.
- Kojima, S. and Mizutani, S. 1987. Triassic and Jurassic Radiolaria from the Nadanhada Range, northeast China. *Transactions and Proceedings of the Palaeontological Society of Japan, New Series*, no. 148, p. 256–275.
- Kojima, S., Yayasaka, Y., Hiroi, Y., Matsuoka, A., Sano, H., Sugamori, Y., Suzuki, N., Takemura, S., Tsujimori, T. and Uchino, T., 2016: Pre-Cretaceous accretionary complexes. *In*, Moreno, T., Wallis, S., Kojima, T., Gibbons, W. *eds.*, *The Geology of Japan*, p. 61–100. The Geological Society, London.
- Kozur, H., 1984: New radiolarian taxa from the Triassic and Jurassic. *Geologisch-Paläontologische Mitteilungen Innsbruck*, Band 13, p. 49–88.
- Li, G. and Bengtson, P., 2018: Early Cretaceous ammonites from eastern Heilongjiang, northeastern China, and their chronostratigraphical significance, *Cretaceous Research*, vol. 88, p. 197–209.
- Li, G. and Matsuoka, A., 2015: Searching for a non-marine Jurassic/Cretaceous boundary in northeastern China. *Journal of the Geological Society of Japan*, vol. 121, p. 109–122.
- Li, G. and Yang, Q., 2003: Confirmation of an Early Cretaceous age for the Qihulin Formation in eastern Heilongjiang Province, China: constraints from a new discovery of radiolarians. *Cretaceous Research*, vol. 24, p. 691–696.
- Li, G. and Yu, S. M., 2004: Discovery of agglutinated foraminifers from the

- Longzhaogou Group in eastern Heilongjiang Province. *Science in China Series D, Earth Sciences*, vol. 47, p. 122–127.
- Li, W. K., Han, J. X., Zhang, S. X. and Meng, F. Y., 1979: The main characteristics of the Upper Palaeozoic stratigraphy at the north Nadanhada Range, Heilongjiang Province, China. *Bulletin of Chinese Academy of Geological Sciences*, vol. 1, p. 104–120. (in Chinese with English abstract)
- Li, W. R., 1996: The Jurassic–Cretaceous framework in the offshore basin in the eastern part of Heilongjiang Province. *Heilongjiang Geology*, vol. 7, p. 1–10. (in Chinese with English abstract)
- Matsuoka, A., 1982: Middle and Late Jurassic radiolarian biostratigraphy in the Sakawa and the Niyodo areas, Kochi Prefecture, southwest Japan. In, Nakaseko, K. ed., *Proceedings of the first Japanese Radiolarian Symposium. News of Osaka Micropalaeontologists*, Special Volume no. 5, p. 237–253. (in Japanese with English abstract)
- Matsuoka, A., 1983: Middle and Late Jurassic radiolarian biostratigraphy in the Sakawa and adjacent areas, Shikoku, southwest Japan. *Journal of Geosciences, Osaka City University*, vol. 26, p. 1–48.
- Matsuoka, A., 1984: Late Jurassic four-segmented Nassellarians (Radiolorai) from Shihoku, Japan. *Journal of Geosciences, Osaka City University*, vol. 27, p. 143–153.
- Matsuoka, A., 1985: Middle Jurassic Keta Foramtion of the southern part of the Middle Chichibu Terrane in the Sakawa area, Kochi Prefecture, southwest Japan. *Journal of the Geological Society of Japan*, vol. 91, p. 411–420. (in Japanese)

with English abstract)

Matsuoka, A., 1986a: *Tricolocapsa yaoi* assemblage (Late Jurassic radiolarians) from the Togano Group in Shikoku, southwest Japan. *Journal of Geosciences, Osaka City University*, vol. 29, p. 101–115.

Matsuoka, A., 1986b: Faunal change of radiolarians around the Jurassic–Cretaceous boundary: with special reference to some multi-segmented nassellarians. *Fossils (Palaeontological Society of Japan)*, no. 40, p. 1–15. (*in Japanese with English abstract*)

Matsuoka, A., 1988: First appearance biohorizon of *Tricolocapsa conexa* within Jurassic siliceous mudstone sequences of the Kamiaso area in the Mino Terrane, central Japan; a correlation of radiolarian zones of the Middle Jurassic. *Journal of the Geological Society of Japan*, vol. 94, p. 583–590.

Matsuoka, A., 1992: Jurassic and Early Cretaceous radiolarians from Leg 129, Sites 800 and 801, western Pacific Ocean. In, Larson, R. L., Lancelot, Y. *et al.*, *Proceedings of the Ocean Drilling Program, Scientific Results, College Station, TX (Ocean Drilling Program)*, vol. 129, p. 203–220.

Matsuoka, A., 1995: Jurassic and Lower Cretaceous radiolarian zonation in Japan and in the western Pacific. *Island Arc*, vol. 4, p. 140–153.

Matsuoka, A., 2006: Revised radiolarian zonation for the entire Jurassic in the western Pacific and Japan. *Volumina Jurassica*, vol. 4, p. 183–184.

Matsuoka, A. and Yao, A., 1985: Latest Jurassic Radiolarians from the Torinosu Group in Southwest Japan. *Journal of Geosciences, Osaka City University*, vol.

28, p. 125–145.

Matsuoka, A. and Yao, A., 1986: A newly proposed radiolarian zonation for the Jurassic of Japan. *Marine Micropaleontology*, vol. 11, p. 91–106.

Matsuoka, A., Yang, Q., Kobayashi, K., Takei, M., Nagahashi, T., Zeng, Q. G. and Wang, Y. J., 2002: Jurassic–Cretaceous radiolarian biostratigraphy and sedimentary environments of the Ceno-Tethys: records from the Xialu Chert in the Yarlung-Zhangbo Suture Zone, southern Tibet. *Journal of Asian Earth Sciences*, vol. 20, p. 277–278.

Matsuoka, A., Yang, Q. and Takei, M., 2005: Latest Jurassic–earliest Cretaceous radiolarian fauna from the Xialu chert in the Yarlung Zangbo suture zone, southern Tibet: Comparison with coeval western Pacific radiolarian faunas and paleoceanographic implications. *Island Arc*, vol. 14, p. 338–345.

Mizutani, S., 1981: A Jurassic formation in the Hida-Kanayama area, central Japan. *Bulletin of the Mizunami Fossil Museum*, no. 8, p. 147–190. (in Japanese with English abstract and description)

Mizutani, S., Kojima, S., Shao, J. A. and Zhang, Q. L., 1986: Mesozoic radiolarians from the Nadanhada area, Northeast China. *Proceedings of the Japan Academy, Series B*, vol. 62, p. 337–340.

Nishihara, C. and Yao, A., 2005: Faunal change of Middle Jurassic (Bajocian) radiolarians in the Inuyama area of the Mino Terrane. *Fossils (Palaeontological Society of Japan)*, no. 78, p. 32–39. (in Japanese with English abstract)

O'Dogherty, L., 1994. Biochronology and paleontology of Mid-Cretaceous

radiolarian from Northern Apennines (Italy) and Betic Cordillera (Spain).

Mémoires de Géologie, Lausanne, vol. 21, p. 1–415.

O'Dogherty, L., Bill, M., Goričan, Š., Dumitrica, P. and Masson, H., 2005: Bathonian

Radiolarians from an Ophiolitic Mélange of the Alpine Tethys (Gets Nappe,

Swiss-French Alps). *Micropaleontology*, vol. 51, p. 425–485.

O'Dogherty, L., Goričan, Š. and Gawlick, H. J., 2017: Middle and Late Jurassic

radiolarians from the Neotethys suture in the Eastern Alps. *Journal of*

Paleontology, vol. 91, p. 25–72.

Pessagno, E. A., Jr., 1971: Jurassic and Cretaceous Hagiastriidae from the

Blake-Bahama Basin (site 5A, JOIDES Leg 1) and the Great Valley sequence,

California Coast Ranges. *Bulletin of American Paleontology*, vol. 60, p. 5–83.

Pessagno, E. A., Jr., 1973: Upper Cretaceous Spumellariina from the Great Valley

Sequence, California Coast Ranges. *Bulletin of American Paleontology*, vol. 63, p.

45–102, pl. 1–21.

Pessagno, E. A., Jr., 1976: Radiolarian zonation and stratigraphy of the Upper

Cretaceous portion of the Great Valley Sequence, California Coast Ranges.

Micropaleontology, Special Publication, vol. 2, p. 1–95, pl. 1–14.

Pessagno, E. A., Jr., 1977a: Upper Jurassic Radiolaria and radiolarian biostratigraphy

of the California Coast Ranges. *Micropaleontology*, vol. 23, p. 56–113.

Pessagno, E. A., Jr., 1977b: Lower Cretaceous radiolarian biostratigraphy of the

Great Valley Sequence and Franciscan Complex, California Coast Ranges.

Cushman Foundation For Foraminiferal Research Special Publication, vol. 15, p.

1–87, pl. 1–12.

Pessagno, E. A., Jr. and Newport, R. L., 1972: A Technique for Extracting Radiolaria from Radiolarian Cherts. *Micropaleontology*, vol. 18, p. 231–234.

Pessagno, E. A., Jr. and Whalen, P. A., 1982: Lower and Middle Jurassic Radiolaria (multicyrtids Nassellariina) from California, east-central Oregon, and the Queen Charlotte Islands, British Columbia. *Micropaleontology*, vol. 28, p. 111–169.

Pessagno, E. A., Jr., Six, W. M. and Yang, Q., 1989: Xiphostylidae Haeckel and Paarvivaccidae, n. fam., (Radiolaria) from the Jurassic of North America. *Micropaleontology*, vol. 35, p. 193–255.

Pessagno E. A., Jr, Blome, C. D., Hull, D. M. and Six, W. M., 1993: Jurassic Radiolaria from the Josephine ophiolite and overlying strata, Smith River subterrane (Klamath Mountains), northwestern California and southwestern Oregon. *Micropaleontology*, vol.39, p. 93–166.

Riedel, W. R. and Sanfilippo, A., 1974: Radiolaria from the southern Indian Ocean, DSDP Leg 26. In, Davies, T. A., Luyendyk, B. P. *et al.*, *Initial Reports of the Deep Sea Drilling Project*, vol. 26, p. 771–814. U.S. Government Printing Office, Washington, D.C.

Rüst, D., 1885: Beitrage zur Kenntniss der Fossilen Radilarien aus Gesteinen des Jura. *Palaeontographica*, Band 31, p. 269–321.

Schaaf, A., 1981: Late Early Cretaceous Radiolaria from Deep Sea Drilling Project Leg 62. In, Thiede, J., Vallier, T. L., Dean, W. E., *Initial Reports of Deep Sea Drilling Project*, vol. 62, p. 419–470. U.S. Government Printing Office,

Washington D.C.

Sha, J. G., 1990: Discovery of *Aucellina* (Bivalvia) in eastern Heilongjiang. *Journal of Stratigraphy*, vol. 14, p. 226–230. (in Chinese with English abstract)

Sha, J. G., 2002: Major achievements in studying the Early Cretaceous biostratigraphy of eastern Heilongjiang. *Earth Science Frontiers*, vol. 9, p. 95–101. (in Chinese with English abstract)

Sha, J. G., 2007: Cretaceous stratigraphy of northeast China: non-marine and marine correlation. *Cretaceous Research*, vol. 28, p. 146–170.

Sha, J. G. and Fürsich, F. T., 1993: Biostratigraphy of the Upper Jurassic–Lower Cretaceous bivalves *Buchia* and *Aucellina* of eastern Heilongjiang, northeast China. *Geological Magazine*, vol. 130, p. 533–542.

Sha, J. G., Fürsich, F. T., Grant-Mackie, J. A., 1994: A revised Early Cretaceous age for the Longzhaogou and Jixi groups of eastern Heilongjiang, China, previously considered Jurassic: Paleogeographic implications. *Newsletters on Stratigraphy*, vol. 31, p. 101–114.

Sha, J. G., Hirano, H., Yao, X. G., Pan, Y. H., 2008: Late Mesozoic transgressions of eastern Heilongjiang and their significance in tectonics, and coal and oil accumulation in northeast China. *Palaeogeography, Palaeoclimatology, Palaeoecology*, vol. 263, 119–130.

Sha, J. G., Wang, J. P., Kirillova, G., Pan, Y. H., Cai, H. W., Wang, Y. Q., Yao, X. G., Peng, B., 2009: Upper Jurassic and lower cretaceous of Sanjiang-Middle Amur basin: Non-marine and marine correlation. *Science in China Series D, Earth Sciences*, vol. 52, p. 1873–1889.

- Sun, G., Sha, J. G., Wang, Y. G., Yuan, F. S., Xiong, X. Z. and Piao, T. Y., 1989. Jurassic–Cretaceous boundary of Dong’anzhen, Raohe, Heilongjiang with a new knowledge of *Buchia*. *Acta Palaeontologica Sinica*, vol. 28, p. 579–597. (in Chinese with English abstract)
- Suzuki, H. and Gawlick, H. J., 2009: Jurassic radiolarians from cherty limestones below the Hallstatt salt mine (Northern Calcareous Alps, Austria). *Neues Jahrbuch Für Geologie Und Palaontologie-Abhandlungen*, vol. 251, p. 155–197.
- Takemura, A., 1986: Classification of Jurassic Nassellarians (Radiolaria). *Palaeontographica, Abteilung A, Palaeozoologie-Stratigraphie*, Band 195, p. 29–74.
- Tan, S. H., 1927: Over de damenstelling en het ontstaan van krijt- en mergel-gesteenten van de Molukker. *Jaarboek van het mijnwezen in Nederlandsch Oost-Indie, jaargang 55, 1926, verhandelingen, 3rd gedeelte*, p. 5–165.
- Wang, C. Y., Kang, B. X. and Zhang, H. R., 1986: A discovery of Triassic conodonts in the Nadanhada Range and geological significance. In, Compiling Committee of Plate Tectonics in Northern China ed., *Contributions to the Project of Plate Tectonics in Northern China*, p. 203–214. Geological Publishing House, Beijing. (in Chinese with English abstract).
- Wang, X. Z., 1959: Marine Mesozoic strata of the Raohe Mesozoic folded belt in northeastern China. *Chinese Journal of Geology*, vol. 2, p. 50–51. (in Chinese).
- Wilde, S. A. and Zhou, J. B., 2015: The late Paleozoic to Mesozoic evolution of the eastern margin of the Central Asian Orogenic Belt in China. *Journal of Asian Earth Sciences*, vol. 113, p. 909–921.

- Wu, F. Y., Yang, J. H., Lo, C. H., Wilde, S. A., Sun, D. Y. and Jahn, B. M., 2007: The Heilongjiang Group: a Jurassic accretionary complex in the Jiamusi Massif at the western Pacific margin of northeastern China. *Island Arc*, vol. 16, 156–172.
- Wu, H. R., 1993: Upper Jurassic and Lower Cretaceous radiolarians of Xialu chert, Yarlung Zangbo ophiolite belt, southern Tibet. *Micropaleontology, Special publication*, vol. 6, p. 115–136.
- Yabe, S. and Ohki, K., 1957: *Geologic Map (1: 250,000) of T'ung-Chiang (NL-53-1) with geologic column and brief description*. Compilation Committee, Geological and Mineralogical Resources Far East, Tokyo Geographic Society, Tokyo.
- Yang, Q., 1993: Taxonomic studies of Upper Jurassic (Tithonian) Radiolaria from the Taman Formation, east-central Mexico. *Palaeoworld, Special Issue*, vol. 3, p. 1–164, pl. 1–26.
- Yang, Q. and Matsuoka, A., 1997: A comparative study on Upper Jurassic radiolarian biostratigraphy of the Taman Formation, east-central Mexico and the ODP Site 801B Section, west Pacific. *Science Report of the Niigata University, Series E, (Geology)*, no. 12, p. 29–49.
- Yang, Q., Mizutani, S. and Nagai, H., 1993: Biostratigraphic Correlation between the Nadanhada Terrane of NE China and the Mino Terrane of Central Japan. *Journal of Earth and Planetary Sciences, Nagoya University*, vol. 40, p. 27–43.
- Yao, A., 1979: Radiolarian fauna from the Mino Belt in the northern part of the Inuyama area, Central Japan. Part II: Nassellaria 1. *Journal of Geosciences, Osaka City University*, vol. 22, p. 21–72.
- Zhang, Q. L., Mizutani, S. and Kojima, S., 1997: Radiolaria and correlation study of

terranes. *Acta Palaeontologica Sinica*, vol. 36, p. 245–252.

Zhou, J. B., Wilde, S. A., Zhang, X. Z., Zhao, G. C., Zheng, C. Q., Wang, Y. J. and

Zhang, X. H., 2009: The onset of Pacific margin accretion in NE China:

Evidence from the Heilongjiang high-pressure metamorphic belt. *Tectonophysics*,

vol. 478, p. 230–246.

Zhou, J. B., Cao, J. L., Wilde, S. A., Zhao, G. C., Zhang, J. J. and Wang, B., 2014:

Paleo-Pacific subduction-accretion: Evidence from Geochemical and U-Pb

zircon dating of the Nadanhada accretionary complex, NE China. *Tectonics*, vol.

33, p. 2444–2466.

Zügel, P., 1997: Discovery of a radiolarian fauna from the Tithonian of the Solnhofen

area (Southern Franconian Alb, southern Germany). *Paläontologische Zeitschrift*,

vol. 71, p. 197–209.

Figure and table captions

Figure 1. Schetch map showing the sample locality (P1: dark aster) in the Nadanhada range area, northeastern China.

Figure 2. Upper Jurassic and Lower Cretaceous strata in the study area of eastern Heilongjiang, northeastern China (after Sha *et al.*, 2009; Li and Matsuoka, 2015; Li and Bengtson, 2018).

Figure 3. Columnar section of the P1 section with sampling horizons.

Figure 4. Radiolarian zones (after Matsuoka, 1995, 2006; Matsuoka *et al.*, 2002) and the identified biozones for the Nadanhada radiolarian samples.

Figure 5. All specimens are from sample P1-09 in the Dajiashan area of Hulin County, eastern Heilongjiang Province, northeastern China. All scale bars are 50 μm except for further indication. **1**, Cryptothracic Nassellaria, specimen P1-09-4; **2**, *Transhsuum?* sp., specimen P1-9-3; **3,4**, *Unuma gordus* Hull, 1997; 3, specimen P1-9-4; 4, specimen P1-9-7. **5**, *Caneta?* sp. 1, specimen P1-09-22; **6**, *Tethysetta dhimenaensis* (Baumgartner, 1984), specimen P1-12-9; **7,8**, “*Stichocapsa*” *robusta* Matsuoka, 1984; 7, specimen P1-09-24; 8, specimen P1-9-5; **9,10**, “*Stichocapsa*” *convexa* Yao, 1979, specimen P1-09-10-3; 9, side view; 10, aperture view; **11**, “*Stichocapsa*” sp., specimen P1-9-8; **12,13**, *Striatojaponocapsa synconexa* O'Dogherty, Gorican and

Dumitrica in O'Dogherty *et al.*, 2005, specimen P1-9-28; 12, side view; 13, aperture view, shows the circular depression near the aperture, scale bar 10 μm ; **14**, "*Tricolocapsa*" sp., specimen P1-9-20; **15,16**, "*Tricolocapsa*" *tetragona* Matsuoka, 1983, specimen P1-9-6; 15, side view; 16, enlargement of the left part of the specimen of 15, showing the tetragonal pore frames consisting of the longitudinal plicae and the interposed transverse bars, scale bar 10 μm ; **17**, *Archaeocenosphaera* sp., specimen P1-09-25.

Figure 6. All specimens are from sample P1-012 in the Dajiashan area of Hulin County, eastern Heilongjiang Province, northeastern China. All scale bars are 50 μm .

1,2, *Plegmosphaera*? spp.; 1, specimen P1-12-1; 2, specimen P1-12-32; **3,4**, *Triactoma* sp.; 3, specimen P1-12-4; 4, specimen P1-12-8. **5,6**, *Archaeodictyomitra rigida* Pessagno, 1977a; 5, specimen P1-12-26; 6, specimen P1-12-43; **7**, *Cinguloturris carpatica* Dumitrica in Dumitrica and Mello, 1982, specimen P1-12-11; **8**, *Tethysetta dhimenaensis* (Baumgartner, 1984), specimen P1-12-27; **9**, *Solenotryma ichikawai* Matsuoka and Yao, 1985, specimen P1-12-41; **10**, *Pseudoxitus*? sp., specimen P1-12-19; **11**, *Zhamoidellum ovum* Dumitrica, 1970, specimen P1-12-30.

Figure 7. All specimens are from sample P1-013 in the Dajiashan area of Hulin County, eastern Heilongjiang Province, northeastern China. All scale bars are 50 μm .

1,2, *Praeconocaryomma* sp.; 1, specimen P1-13-5; 2, specimen P1-13-14. **3**, *Orbiculiforma* sp., specimen P1-13-7. **4**, *Ristola* sp., specimen P1-13-11.

Figure 8. All specimens are from sample P1-08 in the Dajiashan area of Hulin County, eastern Heilongjiang Province, northeastern China. All scale bars are 50 μm . **1**, *Archaeocenosphaera* sp., specimen P1-8-29; **2**, *Praeconocaryomma?* sp., specimen P1-08-17; **3**, *Archaeospongoprunum* sp. 1, specimen P1-08-24. **4**, *Archaeospongoprunum* sp. 2, specimen 108x2-08. **5,6**, *Emiluvia praemyogii* Baumgartner, 1984; 5, specimen 108x2-20; 6, specimen P1-08-28; **7**, *Paronaella* sp. 1, specimen 108x2-34, scale bar 100 μm ; **8**, *Paronaella* sp. 2, specimen P1-08-26, scale bar 100 μm ; **9**, *Gongylothorax favosus* Dumitrica, 1970, specimen P1-08-11; **10**, *Eucyrtidiellum ptyctum* (Riedel and Sanfilippo, 1974), specimen 108x2-24; **11**, *Eucyrtidiellum unumaense* (Yao, 1979), specimen P1-8-64; **12,13**, *Archaeodictyomitra rigida* Pessagno, 1977a; 12, specimen P1-8-38; 13, specimen P1-8-39; **14**, *Archaeodictyomitra* sp., specimen P1-08-20; **15**, *Loopus?* sp. 1, specimen P1-08-2; **16**, *Unuma gordus* Hull, 1997, specimen P1-8-69. **17,18**, *Protunuma japonicus* Matsuoka and Yao, 1985; 17, specimen P1-8-1; 18, specimen P1-8-25. **19**, *Cinguloturris fusiformis* Hori, 1999, specimen P1-08-4. **20,21**, *Cinguloturris carpatica* Dumitrica in Dumitrica and Mello, 1982; 20, specimen 8-9; 21, specimen P1-8-46; **22-25**, *Tethysetta dhimenaensis* (Baumgartner, 1984); 22, specimen 108x2-28; 23, specimen P1-8-27; 24, specimen P1-8-37; 25, specimen P1-8-42.

Figure 9. All specimens are from sample P1-08 in the Dajiashan area of Hulin County,

eastern Heilongjiang Province, northeastern China. All scale bars are 50 μm . **1**, *Caneta* sp. 2, specimen P108-1; **2**, *Olanda?* sp., specimen 108x2-25. **3–5**, *Loopus* sp. 2; 3, specimen P1-08-30; 4, specimen 108x2-11; 5, specimen 108x2-12; **6,7**, *Loopus?* sp. 3; 6, specimen P1-08-6; 7, specimen P1-08-25; **8,9**, *Zhamoidellum ovum* Dumitrica, 1970; 8, specimen P1-8-32; 9, specimen P1-8-50; **10**, *Williriedellum marcucciae* (Cortese, 1993), specimen P1-8-40; **11**, *Zhamoidellum* sp. specimen P1-8-44; **12–15**, *Crococapsa hexagona* (Hori, 1999); 12, specimen P1-8-19; 13, specimen P1-8-10; 14, specimen P1-8-15. 15, specimen P1-8-60; **16,17**, *Arcanicapsa funatoensis* (Aita, 1987); 16, specimen P1-8-49; 17, specimen P1-8-59; **18**, *Theocapsomella himedaruma* (Aita, 1987), specimen P1-8-54.

Table 1. Occurrence of radiolarians in four collected samples (P1-09, 08, 012, 013) from the Dalingqiao Formation in eastern Heilongjiang, northeastern China.

Figure 1




Figure 2

	Raohe	Dajiashan, Hulin	Yunshan, Hulin	Jixi, Boli, Muling		
Albian		Dajiashan Group	Baiheshan Fm	Zhushan Fm	Muling Fm	
Aptian			Dajiashan Fm	Yunshan Fm	Jixi Group	Chengzihe Fm
Barremian			Nandatashan Fm	Qihulin Fm		Didao Fm
Haut.				? ↓		
Tith.-Valang.	? ↑					
	Dong'anzen Fm					
	? ↓					
Kimm.		Dalingqiao Fm				

Figure 3

P1 section




Figure 4

		Stages	Code and biozones	Identified biozones
Cretaceous	Lower (part)	Aptian	<i>Turbocapsula costata</i> Z.	<i>Cinguloturris carpatica</i> assemblage Z. <i>Striatojaponocapsa synconexa</i> - "Tricolocapsa" tetragona assemblage Z.
		Barremian	KR3 <i>Aurisatumalis carinatus</i> Z.	
		Hauterivian	KR2 <i>Cecrops septemporatus</i> Z.	
		Valanginian		
		Barriasian	KR1 <i>Pseudodictyomitra carpatica</i> Z.	
Jurassic	Upper	Tithonian	JR8 <i>Loopus primitivus</i> Z.	
		Kimmeridgian	JR7 <i>Hsuum maxwelli</i> Z.	
		Oxfordian	JR6 <i>Kilinora spiralis</i> Z.	
	Middle	Callovian	JR5 <i>Striatojaponocapsa conexa</i> Z.	
		Bathonian		
		Bajocian	JR4 <i>Striatojaponocapsa plicarum</i> Z.	
		Aalenian	JR3 <i>Laxtorum? jurassicum</i> Z.	
	Lower	Toarcian	JR2 <i>Trillus elkhornensis</i> Z.	
		Pliensbachian		
		Sinemurian	JR1 <i>Parahsuum simplum</i> Z.	
Hettangian		JR0 <i>Bipedis horiae</i> Z.		
Tr.	Up.	Rhaetian		

Figure 5




Figure 6




Figure 7




Figure 8




Figure 9




Table 1

	P1-09	P1-08	P1-012	P1-013
<i>Archaeocenosphaera</i> sp.	X	X		
<i>Caneta</i> ? sp. 1	X			
" <i>Stichocapsa</i> " <i>convexa</i> Yao, 1979	X			
" <i>Stichocapsa</i> " <i>robusta</i> Matsuoka, 1984	X			
" <i>Stichocapsa</i> " sp.	X			
<i>Striatojaponocapsa synconexa</i> O'Dogherty et al., 2005	X			
<i>Tethyssetta dhimenaensis</i> (Baumgartner, 1984)	X		X	
<i>Transsuum</i> ? sp.	X			
" <i>Tricolocapsa</i> " <i>tetragona</i> Matsuoka, 1983	X			
" <i>Tricolocapsa</i> " sp.	X			
<i>Unuma gordus</i> Hull, 1997	X			
<i>Arcanicapsa funatoensis</i> (Aita, 1987)		X		
<i>Archaeospongoprunum</i> sp. 1		X		
<i>Archaeospongoprunum</i> sp. 2		X		
<i>Archaeodictyomitra rigida</i> Pessagno, 1977a		X	X	
<i>Archaeodictyomitra</i> sp.		X		
<i>Caneta</i> ? sp. 2		X		
<i>Cinguloturris carpatica</i> Dumitrica in Dumitrica and Mello, 1982		X	X	
<i>Cinguloturris fusiformis</i> Hori, 1999		X		
<i>Crococapsa hexagona</i> (Hori, 1999)		X		
<i>Emiluvia praemyogii</i> Baumgartner, 1984		X		
<i>Eucyrtidiellum ptyctum</i> (Riedel and Sanfilippo, 1974)		X		
<i>Eucyrtidiellum unumaense</i> (Yao, 1979)		X		
<i>Gongylothorax favosus</i> Dumitrica, 1970		X		
<i>Loopus</i> sp. 1		X		
<i>Loopus</i> sp. 2		X		
<i>Loopus</i> sp. 3		X		
<i>Olanda</i> ? sp.		X		
<i>Paronaella</i> sp. 1		X		
<i>Paronaella</i> sp. 2		X		
<i>Praeconocaryomma</i> ? sp.		X		
<i>Protunuma japonicus</i> Matsuoka and Yao, 1985		X		
<i>Theocapsomella himedaruma</i> (Aita, 1987)		X		
<i>Willriedellum marcucciae</i> (Cortese, 1993)		X		
<i>Zhamoidellum</i> sp.		X		
<i>Zhamoidellum ovum</i> Dumitrica, 1970		X	X	
<i>Plegnosphaera</i> ? spp.			X	
<i>Pseudoxitus</i> ? sp.			X	
<i>Solenotryma ichikawai</i> Matsuoka and Yao, 1985			X	
<i>Triactoma</i> sp.			X	
<i>Orbiculiforma</i> sp.				X
<i>Praeconocaryomma</i> sp.				X
<i>Ristola</i> sp.				X