# Paleontological Research



## Papers in Press

"Papers in Press" includes peer-reviewed, accepted manuscripts of research articles, reviews, and short notes to be published in Paleontological Research. They have not yet been copy edited and/or formatted in the publication style of *Paleontological Research*. As soon as they are printed, they will be removed from this website. Please note they can be cited using the year of online publication and the DOI, as follows:

Humblet, M. and Iryu, Y. 2014: Pleistocene coral assemblages on Irabu-jima, South Ryukyu Islands, Japan. *Paleontological Research*, doi: 10.2517/2014PR020.

#### doi:10.2517/2016PR034

Listriodontine suid and tragulid artiodactyls (Mammalia) from the upper Miocene Nakali Formation, Kenya

TAKEHISA TSUBAMOTO<sup>1</sup>, YUTAKA KUNIMATSU<sup>2</sup>, TETSUYA SAKAI<sup>3</sup>, MOTOTAKA SANEYOSHI<sup>4</sup>, DAISUKE SHIMIZU<sup>5</sup>, NAOKI MORIMOTO<sup>5</sup>, HIDEO NAKAYA<sup>6</sup> AND MASATO NAKATSUKASA<sup>5</sup>

<sup>1</sup>Graduate School of Science and Engineering, Ehime University, 2-5 Bunkyo-cho, Matsuyama 790-8577, Japan (e-mail: tsubamoto@sci.ehime-u.ac.jp)
<sup>2</sup>Faculty of Business Administration, Ryukoku University, 67 Tsukamoto-cho, Fukakusa Fushimi-ku, Kyoto 612-8577, Japan
<sup>3</sup>Interdisciplinary Faculty of Science and Engineering, Shimane University, 1060
Nishikawatsu-cho, Matsue 690-8504, Japan
<sup>4</sup>Faculty of Biosphere-Geosphere Science, Okayama University of Science, 1-1
Ridai-cho, Kita-ku, Okayama 700-0005, Japan
<sup>5</sup>Graduate School of Science, Kyoto University, Kitashirakawa Oiwake-cho, Sakyo-ku, Kyoto 606-8502, Japan
<sup>6</sup>Faculty of Science, Kagoshima University, Korimoto 1-21-35, Kagoshima 890-0065,

Japan

Abstract. Two rare artiodactyl mammals from the basal upper Miocene Nakali Formation (ca. 10 Ma) of central Kenya are described. They are cf. Listriodon sp. (Suidae, Listriodontinae) and Dorcatherium cf. pigotti (Ruminantia, Tragulidae), which are the first discoveries of a listriodontine and a tragulid in the formation. Cf. Listriodon sp. is represented by a talonid of a lower molar that has a strongly lophodont hypolophid. Although this listriodontine material is fragmentary, it is comparable in morphology and size to large and fully lophodont species of the genus Listriodon, such as L. splendens and L. pentapotamiae theobaldi. If the Nakali specimen proves to be phyletically closely related to these two species, it indicates that a highly derived lineage of *Listriodon* existed in East Africa around 10 Ma, implying a possible migration of this lineage from Europe/Asia to East Africa during the middle or earliest late Miocene. Dorcatherium cf. pigotti is represented by DP4 (or M1) and a mandible with p3-m3, which are comparable in size to those of *D. pigotti* among the African species of the genus. Although the genus Dorcatherium and species D. pigotti are common taxa in the early to middle Miocene of Africa, they are rarely found in the late Miocene. This is the second record of the genus in the late Miocene of Africa, reinforcing evidence that Dorcatherium existed until the basal late Miocene in East Africa.

Key words: Africa, Dorcatherium, Listriodon, Ruminantia, Tortonian, Vallesian

#### Introduction

The upper Miocene Nakali Formation of central Kenya is an important fossil-bearing deposit particularly for anthropology and primatology because the formation has yielded several taxa of colobine, small non-cercopithecoid catarrhine, and hominoid primates (Nakatsukasa *et al.*, 2006, 2010; Kunimatsu *et al.*, 2007, 2016). Among these fossil primates, a new great ape, *Nakalipithecus nakayamai*, attracted much attention because it is a candidate for the latest common ancestor of the extant African great apes and humans (Kunimatsu *et al.*, 2007). The fossil mammals including primates, a rodent, carnivores, proboscideans, perissodactyls, and artiodactyls from the formation were first reported by Aguirre and Leakey (1974). Since then, additional fossil mammalian specimens have been described and mentioned from the formation by several researchers (Aguirre and Guérin, 1974; Pickford, 1983, 1986; Flynn and Sabatier, 1984; Fischer, 1986; Morales and Pickford, 2006). However, the Nakali fossil mammal fauna has still not been well documented.

Since 2002, a joint research team from Kyoto University and the National Museums of Kenya has been conducting geological and paleontological field research in the Nakali Formation (Nakatsukasa *et al.*, 2006; Nakatsukasa, 2009; Sakai *et al.*, 2013) and has described some mammalian fossil specimens from the formation (Kunimatsu *et al.*, 2007, 2016; Nakatsukasa *et al.*, 2010; Handa *et al.*, 2015; Tsubamoto *et al.*, 2015). This article describes two rare artiodactyls from the Nakali Formation, a listriodontine suid and a tragulid ruminant, collected by the joint research team. These two artiodactyl taxa are rare in terms of collection size in the formation and are important for clarifying the Nakali fossil mammal fauna. *Repository.*—The specimens described here are stored in the Palaeontology Section, National Museums of Kenya, Nairobi, Kenya.

*Taxonomy and terminology.*—Generic taxonomy of the Listriodontinae generally follows van der Made (1996). Dental terminology for the Tragulidae generally follows Sánchez *et al.* (2015). The trigonid stands for a homologous part to the triangle formed by the protoconid, metaconid, and paraconid in the lower molars of primitive tribosphenic mammals; and the talonid stands for a homologous part to the dental crown distal to the trigonid in the lower molars of primitive tribosphenic mammals (Hershkovitz, 1971; Kielan-Jaworowska *et al.*, 2004).

*Dental abbreviations.*—**M**/**m**, upper/lower molars; **P**/**p**, upper/lower premolars; **DP**, upper deciduous premolars.

*Institutional abbreviations.*—**KNM**, National Museums of Kenya, Nairobi, Kenya; **KU**, Kyoto University, Kyoto, Japan; **NMNS-M**, National Museum of Nature and Science, Tsukuba, Japan.

*Other abbreviations.*—**ELMA**, European Land Mammal Age; **MN**, European Neogene Mammal Chron; **NA**, Nakali fossil locality, central Kenya.

#### Locality, formation, and age

The specimens described here are from the basal upper Miocene Nakali Formation at the Nakali locality (approx. 01°11′ N and 36°22′ E), which is located on the eastern margin of the central Kenya Rift, about 40 km west of Maralal, central Kenya (Figure 1; Kunimatsu *et al.*, 2007; Sakai *et al.*, 2013). It is underlain by the Alengerr Tuffs (Golden, 1978), and is unconformably overlain by the Nasorut Formation, which is composed of basalt and trachyte lava, pyroclastics, and conglomerate (Figure 2).

The Nakali Formation (ca. 340 m in thickness) is stratigraphically subdivided into the "Lower," "Middle," and "Upper" members, and is characterized by tuffaceous sandstones and mudstones with conglomerate interbeds, a thick pyroclastic flow deposit, and tuff beds (Figure 2; Kunimatsu *et al.*, 2007; Sakai *et al.*, 2013). The basal part of the "Lower" Member consists mainly of lacustrine turbidite and debrite of lake delta front origin, grading upwards into fluvio-lacustrine deposits. The "Middle" Member consists of a thick pyroclastic flow deposit. The lower half of the "Upper" Member consists mainly of fluvio-lacustrine deposits and the upper half is represented by muddy sediments of lake slope origin (Sakai *et al.*, 2013).

The age of the mammalian and other vertebrate fossils from the Nakali Formation is estimated to be around 10.0-9.8 Ma, that is, early Tortonian (basal late Miocene), based on the anorthoclase  ${}^{40}$ Ar- ${}^{39}$ Ar datings and magnetostratigraphy (Figure 2; Kunimatsu *et al.*, 2007). It ranges within the period from Chron C5n.2n (11.06–9.98 Ma) to C5n.1n (9.94–9.79 Ma), with Chron C5n.1r (9.98–9.94 Ma) corresponding to the horizon from the upper part of the "Lower" Member to the basal part of the "Upper" Member (Kunimatsu *et al.*, 2007; Hilgen *et al.*, 2012; Tsubamoto *et al.*, 2015). Therefore, the Nakali vertebrate fossil fauna is correlated to the uppermost part of MN9 or lowermost part of MN10, and to the Vallesian ELMA (McKenna and Bell, 1997; Hilgen *et al.*, 2012).

#### Systematic paleontology

Order Artiodactyla Owen, 1848

Family Suidae Gray, 1821

Subfamily Listriodontinae Gervais, 1859 Genus *Listriodon* von Meyer, 1846

Type species.—Listriodon splendens von Meyer, 1846.

cf. Listriodon sp.

Figure 3

*Material.*—KNM-NA 66161, a talonid fragment of a left lower molar.

Sublocality.—KU Sublocality Number NA22.

*Horizon.*—"Lower" Member of the Nakali Formation (Figure 2).

*Measurement.*—Talonid (or distal lophid in the Suidae) width = 17.6 mm.

*Description.*—This lower molar talonid (or distal lophid in the Suidae) is lophodont and is moderately worn. The mesial and distal margins are broken (Figure 3F, G), and the root is poorly preserved (Figure 3E). The hypoconid and entoconid are sharp and are connected to each other by the buccolingually oriented hypolophid. The hypolophid is well developed and high for the family. The entoconid is mesiodistally compressed in lingual view (Figure 3C). The entoconid is higher than the hypoconid at this wear stage: the hypoconid is more worn than the entoconid. The prehypocristid (= cristid obliqua) is weak and originates at the mesial face of the hypolophid, extending mesiolingually. It is much more lingually located compared to the preserved tip of the hypoconid. There is no cristid on the distal face of the hypolophid. There is a small cuspule at the distal base of the hypoconid. No cingulid is observed. The root for the talonid is buccolingually bifurcate (Figure 3E–G). The enamel is not crenulated.

Comparisons.—KNM-NA 66161 is assigned to the Listriodontinae, which are bilophodont suids. It has a lophodont hypolophid and a relatively sharp and high entoconid. This morphology and size are reminiscent of the lower molars of some bilophodont suoid artiodactyls (such as the listriodontine suids and some palaeochoerids), tapirid perissodactyls, and macropodid marsupials. KNM-NA 66161 is distinguished from the molariform teeth of the pyrotheriids (South American mammals) and Neogene proboscideans, which have lophodont teeth, in being much smaller and in having much thinner enamel and a cristid/crista extending diagonally from the buccolingually oriented lophid/loph. It is distinguished from the upper molars of the above mentioned mammals in having much less buccolingually separated roots. It differs from the lower molars of the tapirids in having a much more lingually shifted prehypocristid and a much more mesiodistally compressed entoconid. It also differs from the lower molars of the macropodids in having a much more mesiodistally compressed entoconid and a much more diagonally oriented prehypocristid at the central position. Furthermore, it differs from the lower molars of some palaeochoerids that have a lophodont dentition such as *Schizoporcus* and *Yunnanochoerus* in having a much more lingually shifted prehypocristid, a much more mesiodistally compressed entoconid, and more lophodont teeth with a lack of a deep V-shaped notch on the hypolophid in distal view. Therefore, it is identified as a fragmentary lower molar of the Listriodontinae.

KNM-NA 66161 is comparable in morphology to the lower molars of *Listriodon* and *Lopholistriodon*, which have a very derived degree of lophodonty among the subfamily. It differs from the lower molars of the other listriodontine genera such as

*Nguruwe, Kubanochoerus, Bunolistriodon*, and *Eurolistriodon* in having a much more lophodont dentition (van der Made, 1996; Orliac, 2009; Orliac *et al.*, 2009). *Listriodon* and *Lopholistriodon* are the most lophodont and derived genera among the subfamily. These two genera are distinguished from each other mainly based on size and morphology of the upper canines (van der Made, 1996). The upper canine morphology cannot apply to the identification of KNM-NA 66161 because this specimen lacks upper canines. The species of *Listriodon* are much larger than those of *Lopholistriodon*.

KNM-NA 66161 is likely assigned to *Listriodon* rather than to *Lopholistriodon*. It has a talonid width of 17.6 mm, which is comparable in size to m2 of *Listriodon*, is much larger than m1 of *Listriodon* and the lower molars of *Lopholistriodon*, and is smaller than m3 of Listriodon (van der Made, 1996). Even when compared to the species of Listriodon and Lopholistriodon, KNM-NA 66161 has relatively well-developed lophodonty with a mesiodistally compressed entoconid, which is a derived characteristic. In this morphology, it is comparable to the lower molars of the more lophodont species such as Lopholistriodon kidogosana, Listriodon splendens, Listriodon pentapotamiae theobaldi, and Listriodon bartulensis (including 'Listriodon akatikubas' from the Fort Ternan locality of Kenya reported by Pickford, 1986) (van der Made, 1996; Pickford, 2001b). KNM-NA 66161 is larger than the lower molars of Lo. kidogosana and Li. bartulensis, so that it is most comparable to the lower molars of Li. splendens and Li. pentapotamiae theobaldi in morphology and size. Therefore, it is likely assigned to the genus Listriodon. However, a possibility that KNM-NA 66161 may be derived from the lineage of Lopholistriodon with an increased size and therefore may be assigned to the genus *Lopholistriodon* cannot be completely excluded because the present material from the Nakali Formation is poor. KNM-NA 66161 appears to

have a more compressed entoconid than the lower molars of *Li. splendens* and *Li. pentapotamiae theobaldi*. Therefore, this specimen might represent a new species of *Listriodon* (or perhaps *Lopholistriodon*), although it is not confirmed due to the scarcity of the material.

Suborder Ruminantia Scopoli, 1777 Family Tragulidae Milne-Edwards, 1864 Genus *Dorcatherium* Kaup, 1833

*Type species.*—*Dorcatherium naui* Kaup, 1833.

Dorcatherium cf. pigotti Whitworth, 1958

Figures 4–7

*Material.*—KNM-NA 54202, a right mandible with p3–m3; KNM-NA 57182, left

DP4 or M1.

Sublocality.—KU Sublocality Number NA39.

Horizon.—"Upper" Member of the Nakali Formation (Figure 2).

Dental measurements.—Shown in Table 1.

Description.—KNM-NA 57182 is a left upper molariform tooth and is moderately

worn (Figure 4). The crown is low and brachyodont with conical buccal main cusps

(paracone and metacone) and selenodont lingual main cusps (protocone and

metaconule). There are interstitial wear facets on the mesial face of the crown lingual to

the parastyle and on the distal face of the crown near the distal base of the metacone,

indicating that the tooth is not M3, but M1, M2, or DP4. The buccal rib of the paracone is stronger than that of the metacone. The parastyle is located mesially to the paracone. The mesostyle is located distobuccally to the paracone and mesiobuccally to the metacone. The ectoloph connects the parastyle, paracone, mesostyle, metacone, and distobuccal base of the crown. The metastyle is absent. The preprotocrista extends mesiobuccally, connecting to the parastyle. The postprotocrista extends distally and connects to the mesial wall of the metaconule. The premetaconule crista extends mesiobuccally, fading out at the valley between the paracone and the metacone: it does not connect to either the mesostyle or the ectoloph. The postmetaconule crista extends distobuccally and connects to the postmetacrista at the distal base of the metacone (= at the distobuccal corner of the crown). The paraconule is absent. The mesial cingulum exists at the mesial base of the protocone. The lingual cingulum exists between the protocone and the metaconule. The distal cingulum originates at the distal base of the metaconule and is continuous to the lingual base of the metaconule: it does not connect to the lingual cingulum but is located buccally to the lingual cingulum at the lingual base of the metaconule. There is no buccal cingulum. The roots for the paracone and metacone are well separated from each other. The root for the metaconule is smaller than that for the protocone (Figure 4D). These two lingual roots are located close to each other and are fused with each other near the cervix of the tooth.

KNM-NA 54202 is a right mandible with p3–m3 (Figures 5–7). The ventral border of the mandibular corpus is convex and its deepest part is below the trigonid of m3 (Figure 5). The mandibular depth is: 13.7 mm below m2 at the lingual side; 13.5 mm below m2 at the buccal side; 15.4 mm below m3 trigonid at the lingual side; and 14.5 mm below m3 trigonid at the buccal side. The teeth are brachyodont.

The p3 is simple and mesiodistally elongated (Figures 6, 7). Its mesial part is broken. There is a small cusp (= hypoconid?) distal to the main large cusp (= protoconid). There are buccal and lingual grooves between these two cusps. A cristid extends from the protoconid to the distal margin of the crown, connecting the protoconid to the distal cuspid. There is a cingulid at the distolingual base of the distal cuspid. There is no buccal cingulid. Judging from the size and position of the p3 mesial root, p3 appears to be longer than p4.

The p4 is also simple and mesiodistally elongated (Figures 6, 7). It is wider than p3. There is a tall protoconid. The preprotocristid is oriented mesiolingually rather than mesially, bearing a small swelling at the paraconid position. The postprotocristid extends distally, bearing a small cuspule or cristid at the lingual side of the basal position, and connects to the distal margin of the crown, where a low hypoconid appears to occur. The postprotocristid is more heavily worn than the preprotocristid. The lingual postprotocristid appears to be absent. There are mesial and distal cingulids.

The lower molars are basically tetracuspidate with a hypoconulid on m3 (Figure 6). The lower molar size gradient is: m1 < m2 < m3. The buccal cusps (protoconid and hypoconid) are selenodont, and the lingual cusps (metaconid and entoconid) are conical and are somewhat buccolingually compressed. The m1 is heavily worn and m2 is moderately worn. On m1–m2, there are mesial and distal cingulids, lacking buccal and lingual cingulids, and the posthypocristid extends distolingually, fading out at the distal base of the entoconid.

The m3 is also worn, but the precise structure of the cristids can be observed (Figure 6). The preprotocristid and premetacristid are connected to each other at the

mesial margin of the crown. These two cristids do not protrude much mesially and make a Dorcatherium-platform (Sánchez et al., 2015). The postprotocristid and postmetacristid are connected with each other, making a V-shaped notch at the distal trigonid wall. There is a *Dorcatherium*-fold at the distolingual face of the metaconid. A cristid that is located buccally to the postprotocristid exists on the distal face of the protoconid. This cristid appears to originate from the tip of the protoconid and extends distolingually, approaching the mesial tip of the prehypocristid (= cristid obliqua). Therefore, this cristid is identified as a Tragulus-fold, not as a Palaeomeryx-fold (Geraads, 2010). The *Dorcatherium*-fold, postmetacristid, postprotocristid, and Tragulus-fold make an M-structure (Métais et al., 2001) on the distal trigonid wall. The prehypocristid extends mesiolingually and reaches the distal wall of the protoconid. It does not connect to the preentocristid. The posthypocristid extends distolingually. The preentocristid connects to the notch of the distal trigonid wall. The postentocristid is absent or very weak. The hypoconulid is single-cusped with only one cristid, lacking a hypoconulid basin and a hypoconulid loop. The cristid on the hypoconulid extends mesiolingually from the tip of the hypoconulid and connects to the posthypocristid. There is a mesial cingulid. The lingual cingulid only exists at the lingual base of the hypoconulid. There are no buccal and distal cingulids.

*Comparisons.*—The present specimens have a dental morphology typical of small primitive ruminants, such as a brachyodont dentition, a relatively weakly developed selenodonty, no paraconule on the upper molar/DP4, and no molar paraconid. They are assigned to the Tragulidae in having an M-structure with *Dorcatherium-* and *Tragulus*-folds on the distal trigonid wall of the lower molars and morphologically simpler premolars than those of the pecorans, and in lacking a *Palaeomeryx*-fold.

Among the Tragulidae, the present specimens are assigned to Dorcatherium, which is a wastebasket genus in the family, mainly defined as having a *Dorcatherium*-platform on the lower molars. The specimens are distinguished from Dorcabune and *Krabitherium* in having a much less bunodont dentition. They are distinguished from Archaeotragulus in lacking a paralophid on the lower molars, any trace of the hypoconulid on m2, and a Zhailimeryx-fold (= 'entoconidian groove' in Métais et al., 2001) on the mesial face of the molar entoconid. They are distinguished from Afrotragulus and Yunnanotherium in being much larger. They are also distinguished from Afrotragulus, Siamotragulus, Moschiola, Hyemoschus, and Tragulus in having a less mesially protruded preprotocristid and premetacristid with a Dorcatherium-platform on the lower molars. They differ from Afrotragulus in having a cingulum on the metaconule on the upper molar/DP4, a stronger M-structure on the distal trigonid wall on the lower molars, a mesiodistally shorter central valley between the trigonid and the talonid without an interlobular bridge (Sánchez et al., 2010) on the lower molars. They differ from *Moschiola* in having a *Tragulus*-fold. They differ from Hyemoschus in having a distally oriented postprotocrista (distobuccally oriented in *Hyemoschus*) and weaker buccal ribs of the paracone and metacone on the upper molar/DP4. They differ from *Tragulus* in having a stronger mesostyle and parastyle on the upper molar/DP4 and a less buccolingually compressed metaconid and entoconid on the lower molars (Figure 8).

The genus *Dorcatherium* includes many species and is recorded in the Miocene of Africa and Europe and in the Miocene and early Pliocene of Asia (Pickford, 2001a, 2002; Rössner, 2007; Geraads, 2010; see also Sánchez *et al.*, 2010, 2015). The species identification within each continent has been mainly based on the size (Colbert, 1935;

Whitworth, 1958; Pickford, 2001a, 2002). The present specimens are comparable in size to *Dorcatherium pigotti* among the African species of the genus. The length of the lower molar row of KNM-NA 54202 is 29.6 mm, which is comparable to that of *D. pigotti* (Pickford, 2001a, 2002; Geraads, 2010). The size of the upper molariform tooth (KNM-NA 57182) is comparable to or slightly smaller than M1 of this species (Whitworth, 1958), implying that KNM-NA 57182 appears to be M1 or DP4 of the same species as the mandibular specimen KNM-NA 54202. KNM-NA 57182 appears to be DP4 because its mesiobuccal corner projects out somewhat mesially, which is often seen in DP4 but is not seen in the upper molars among the species of *Dorcatherium*.

The mandibular specimen KNM-NA 54202 differs from *D. pigotti* and is unique in the genus *Dorcatherium* in lacking a lingual postprotocristid on p4 and a hypoconulid basin on m3. *D. pigotti* and other species of *Dorcatherium* have a lingual postprotocristid on p4, making a groove on the distal face of the protoconid between the postprotocristid and the lingual postprotocristid (Colbert, 1935; Whitworth, 1958; Ginsburg *et al.*, 2001; Pickford, 2002; Aiglstorfer *et al.*, 2014). Also, *Dorcatherium* usually has a hypoconulid basin with two (mesial and buccal) cristids (or a hypoconulid loop) on m3, whereas KNM-NA 54202 has only a single cristid on the m3 hypoconulid without a hypoconulid basin. We should note that this m3 character might be an individual variation within a species because the same morphological variation is observed in the specimens of a living tragulid species *Tragulus javanicus* stored in NMNS (Figure 8): most specimens have a hypoconulid basin on m3, while one specimen (NMNS M31085) lacks it. Because the material from the Nakali Formation is poor, it is impossible to determine whether these two characteristics indicate intraspecific variation in *D. pigotti* or interspecific differences. Therefore, although

there is a possibility that the Nakali *Dorcatherium* could be a new species based on the morphology of p4 and the m3 hypoconulid, it is not clear due to the scarcity of the fossil material from the Nakali Formation.

#### **Discussion and conclusions**

The two taxa described here (cf. *Listriodon* sp. and *Dorcatherium* cf. *pigotti*) are the first discoveries of a listriodontine and a tragulid in the Nakali Formation, adding two taxa to the faunal list of the formation (Kunimatsu *et al.*, 2007).

The discovery of a listriodontine lower molar (KNM-NA 66161; Figure 3) indicates that a *Listriodon*-like very lophodont and large listriodontine suid existed in the Nakali area at the time of deposition. The genus *Listriodon* (*sensu* van der Made, 1996) has not been recorded in Africa except for a small and highly lophodont species of *Listriodon*, *Li. bartulensis* from the Member A (ca. 13 Ma) of the Ngorora Formation (Tugen Hills) and from the Fort Ternan locality (ca. 14 Ma), Kenya (Pickford, 2001a). If the Nakali listriodontine is phyletically closely related to the large and lophodont species of *Listriodon*, *Li. splendens* (recorded in Europe, Turkey, and East Asia; van der Made, 1996), it indicates that a highly derived lineage of *Listriodon* existed in East Africa around 10 Ma and also implies a possibility that this lineage might have migrated from Europe/Asia to East Africa during the middle or earliest late Miocene.

*Dorcatherium* cf. *pigotti* discovered in the Nakali Formation is the second record of the genus in the late Miocene of Africa (Pickford, 1991; Geraads, 2010). Although the genus *Dorcatherium* and species *D. pigotti* are common taxa in the early to middle

Miocene of Africa, they are rarely found in the late Miocene (Geraads, 2010). In the Pliocene (ca. 5.0 Ma), the living tragulid genus of Africa, *Hyemoschus*, appeared in East Africa (Pickford *et al.*, 2004; Geraads, 2010; *contra* Rössner, 2007). This study reinforces the observation that an archaic tragulid *Dorcatherium* survived until the base of the late Miocene in East Africa before the appearance of the living tragulid genus of Africa, *Hyemoschus*.

### Acknowledgments

We are grateful to the National Commission for Science, Technology and Innovation (NACOSTI), the Government of Kenya for according research permission for us. We thank Fredrick Kyalo Manthi, Margaret Omoto, Mary Muungu, and Francis Ndiritu Muchemi (National Museums of Kenya, Nairobi, Kenya) for kindly providing access to the specimens examined. We are also grateful to Shin-ichiro Kawada (National Museum of Nature and Science, Tsukuba, Japan) for graciously providing access to the comparative specimens. Thanks are also due to Daisuke Mizoguchi and Chiharu Kamimura (Nairobi Research Station, Japan Society for the Promotion of Science, Nairobi, Kenya) for their help during our stay in Kenya. This manuscript was improved with critical reviews by Martin Pickford (Muséum National d'Histoire Naturelle, Paris, France) and Jan van der Made (Museo Nacional de Ciencias Naturales, Madrid, Spain). This research was supported by JSPS KAKENHI Grant Numbers 25257408 and 16H02757 (to M. Nakatsukasa) and 16K07534 (to T. Tsubamoto).

#### References

- Aguirre, E. and Guérin, C., 1974: Première découverte d'un Iranotheriinae (Mammalia, Perissodactyla, Rhinocerotidae) en Afrique: *Kenyatherium bishopi* nov. gen. sp. de la formation vallésienne (Miocene supérieur) de Nakali (Kenya). *Estudios Geológicos*, vol. 30, p. 229–233.
- Aguirre, E. and Leakey, P., 1974: Nakali: nueva fauna de *Hipparion* del Rift Valley de Kenya. *Estudios Geológicos*, vol. 30, p. 219–227.
- Aiglstorfer, M., Rössner, G. E. and Böhme, M., 2014: *Dorcatherium naui* and pecoran ruminants from the late Miocene Gratkorn locality (Austria). *Palaeobiodiversity and Palaeoenvironments*, vol. 94, p. 83–123.
- Colbert, E. H., 1935: Siwalik mammals in the American Museum of Natural History. *Transactions of the American Philosophical Society, Philadelphia, New Series*, vol. 26, p. x + 1–401.
- Fischer, M. S., 1986: Die Stellung der Schliefer (Hyracoidea) im phylogenetischen System der Eutheria. Zugleich ein Beitrag zur Anpassungsgeschichte der Procaviidae. *Courier Forschungsinstitut Senckenberg*, no. 84, p. 1–132.
- Flynn, L. J. and Sabatier, M., 1984: A muroid rodent of Asian affinity from the Miocene of Kenya. *Journal of Vertebrate Paleontology*, vol. 3, p. 160–165.
- Geraads, D., 2010: Tragulidae. In, Werdelin, L. and Sanders, W. J. eds., Cenozoic Mammals of Africa, p. 723–729. University of California Press, Berkeley.
- Gervais, P., 1859: Zoologie et Paléontologie françaises, 2nd Edition, 544 p. Bertrand, Paris.
- Ginsburg, L., Morales, J. and Soria, D., 2001: Les Ruminantia (Artiodactyla, Mammalia) du Miocène des Bugti (Balouchistan, Pakistan). *Estudios Geológicos*,

vol. 57, p. 155–170.

- Golden, M., 1978: *The Geology of the Area East of Silali, Rift Valley Province, Kenya*,254 p. Ph.D. Dissertation, University of London, London.
- Gray, J. E., 1821: On the natural arrangement of vertebrose animals. *London Medical Repository*, vol. 15, p. 296–310.
- Handa, N., Nakatsukasa, M., Kunimatsu, Y., Tsubamoto, T. and Nakaya, H., 2015: New specimens of *Chilotheridium* (Perissodactyla, Rhinocerotidae) from the Upper Miocene Namurungule and Nakali Formations, northern Kenya. *Paleontological Research*, vol. 19, p. 181–194.
- Hershkovitz, P., 1971: Basic crown patterns and cusp homologies of mammalian teeth. *In*, Dahlberg, A. A. *ed.*, *Dental Morphology and Evolution*, p. 95–150. The
  University of Chicago Press, Chicago.
- Hilgen, F. J., Lourens, L. J. and Van Dam, J. A., 2012: The Neogene Period. *In*, Gradstein, F. M., Ogg, J. G., Schmitz, M. D. and Ogg, G. M. *eds.*, *The Geologic Time Scale 2012*, p. 923–978. Elsevier, Amsterdam.
- Kaup, J. J., 1833: Über die Gattung Dinotherium, Zusätze und Verbesserungen zum ersten heft der description d'ossemens fossiles. Neues Jahrbuch für Mineralogie, Geologie und Paläontologie, vol. 1833, p. 509–517, pl. 7.
- Kielan-Jaworowska, Z., Cifelli, R. L. and Luo, Z.-X., 2004: Mammals from the Age of Dinosaurs: Origins, Evolution, and Structure, xv + 630 p. Columbia University Press, New York.
- Kunimatsu, Y., Nakatsukasa, M., Sawada, Y., Sakai, T., Hyodo, M., Hyodo, H., Itaya,T., Nakaya, H., Saegusa, H., Mazurier, A., Saneyoshi, M., Tsujikawa, H.,Yamamoto, A. and Mbua, E., 2007: A new Late Miocene great ape from Kenya and

its implications for the origins of African great apes and humans. *Proceedings of the National Academy of Sciences of the United States of America*, vol. 104, p. 19220–19225.

- Kunimatsu, Y., Nakatsukasa, M., Sawada, Y., Sakai, T., Saneyoshi, M., Nakaya, H., Yamamoto, A. and Mbua, E., 2016: A second hominoid species in the early Late Miocene fauna of Nakali (Kenya). *Anthropological Science*, vol. 124, p. 75–83.
- McKenna, M. C. and Bell, S. K., 1997: *Classification of Mammals, above the Species Level*, xii + 631 p. Columbia University Press, New York.
- Métais, G., Chaimanee, Y., Jaeger, J.-J. and Ducrocq, S., 2001: New remains of primitive ruminants from Thailand: evidence of the early evolution of the Ruminantia in Asia. *Zoologica Scripta*, vol. 30, p. 231–248.
- Milne-Edwards, A., 1864: Recherches anatomiques, zoologiques, et paléontologiques sur la famille des chevrotains. *Annales des Sciences Naturelles*, *Zoologie*, *Paris*, *Ser. 5*, vol. 2, p. 49–167, pls. 2–12.
- Morales, J. and Pickford, M., 2006: A large percrocutid carnivore from the Late Miocene (ca. 10–9 Ma) of Nakali, Kenya. *Annales de Paléontologie*, vol. 92, p. 359–366.
- Nakatsukasa, M., 2009: Paleontological research in Nakali, Kenya. *Anthropological Science (Japanese Series)*, vol. 117, p. 111–117. (*in Japanese*)
- Nakatsukasa, M., Kunimatsu, Y., Sawada, Y., Sakai, T., Hyodo, H., Itaya, T., Saneyoshi,
  M., Tsujikawa, H. and Mbua, E., 2006: Late Miocene primate fauna in Nakali,
  central Kenya. *American Journal of Physical Anthropology*, vol. 129 (S42), p. 136.
- Nakatsukasa, M., Mbua, E., Sawada, Y., Sakai, T., Nakaya, H., Yano, W. and Kunimatsu, Y., 2010: Earliest colobine skeletons from Nakali, Kenya. *American*

Journal of Physical Anthropology, vol. 143, p. 365–382.

- Orliac, M. J., 2009: The differentiation of bunodont Listriodontinae (Mammalia, Suidae) of Africa: new data from Kalodirr and Moruorot, Kenya. *Zoological Journal of the Linnean Society*, vol. 157, p. 652–677.
- Orliac, M. J., Antoine, P.-O., Métais, G., Marivaux, L., Crochet, J.-Y., Welcomme, J.-L.,
  Baqri, S. R. H. and Roohi, G., 2009: *Listriodon guptai* Pilgrim, 1926 (Mammalia,
  Suidae) from the early Miocene of the Bugti Hills, Balochistan, Pakistan: new
  insights into early Listriodontinae evolution and biogeography. *Naturwissenschaften*, vol. 96, p. 911–920.
- Owen, R., 1848: Description of teeth and portions of jaw of two extinct anthracotherioid quadrupeds (*Hyopotamus vectianus* and *Hyop. bovinus*) discovered by the Marchioness of Hastings in the Eocene deposits of the N.W. coast of the Isle of Wight: with an attempt to develop Cuvier's idea of the classification of Pachyderms by the number of their toes. *Quarterly Journal of the Geological Society of London*, vol. 4, p. 103–141, pls. 7, 8.
- Pickford, M., 1983: On the origins of Hippopotamidae together with descriptions of two species, a new genus and a new subfamily from the Miocene of Kenya. *Geobios*, vol. 16, p. 193–217.
- Pickford, M., 1986: A revision of the Miocene Suidae and Tayassuidae, (Artiodactyla, Mammalia) of Africa. *Tertiary Research Special Paper*, no. 7, p. 1–83.
- Pickford, M., 1991: Biostratigraphic correlation of the Middle Miocene mammal locality of Jabal Zaltan, Libya. *In*, Salem, M. J., Hammuda, O. S. and Eliagoubi, B. A. *eds.*, *Third Symposium on the Geology of Libya (Tripoli, 1987)*, p. 1483–1490. Elsevier, Amsterdam.

- Pickford, M., 2001a: Africa's smallest ruminant: A new tragulid from the Miocene of Kenya and the biostratigraphy of East African Tragulidae. *Geobios*, vol. 34, p. 437–447.
- Pickford, M., 2001b: New species of *Listriodon* (Suidae, Mammalia) from Bartule, Member A, Ngorora Formation (ca 13 Ma), Tugen Hills, Kenya. *Annales de Paléontologie*, vol. 87, p. 207–221.
- Pickford, M., 2002: Ruminants from the early Miocene of Napak, Uganda. *Annales de Paléontologie*, vol. 88, p. 85–113.
- Pickford, M., Senut, B. and Mourer-Chauviré, C., 2004: Early Pliocene Tragulidae and peafowls in the Rift Valley, Kenya: Evidence for rainforest in East Africa. *Comptes Rendus Palevol*, vol. 3, p. 179–189.
- Rössner, G. E., 2007: Family Tragulidae. In, Prothero, D. R. and Foss, S. E. eds., The Evolution of Artiodactyls, p. 213–220. Johns Hopkins University Press, Baltimore.
- Sakai, T., Saneyoshi, M., Sawada, Y., Nakatsukasa, M., Kunimatsu, Y. and Mbua, E.,
  2013: Early continental rift basin stratigraphy, depositional facies and tectonics in volcaniclastic system: examples from the Miocene successions along the Japan Sea and in the East African Rift Valley (Kenya). *In*, Itoh, Y. *ed.*, *Mechanism of Sedimentary Basin Formation—Multidisciplinary Approach on Active Plate Margin*, p. 83–108. InTech, Rijeka.
- Sánchez, I. M., Quiralte, V., Morales, J. and Pickford, M., 2010: A new genus of tragulid ruminant from the early Miocene of Kenya. *Acta Palaeontologica Polonica*, vol. 55, p. 177–187.
- Sánchez, I. M., Quiralte, V., Ríos, M., Morales, J. and Pickford, M., 2015: First African record of the Miocene Asian mouse-deer *Siamotragulus* (Mammalia, Ruminantia,

Tragulidae): implications for the phylogeny and evolutionary history of the advanced selenodont tragulids. *Journal of Systematic Palaeontology*, vol. 13, p. 543–556.

- Scopoli, G. A., 1777: Introductio ad historiam naturalem, sistens genera lapidum, plantarum et animalium hactenus detecta, caracteribus essentialibus donata, in tribus divisa, subinde ad leges naturae, 506 p. Prague, Gerle.
- Tsubamoto, T., Kunimatsu, Y., Nakaya, H., Sakai, T., Saneyoshi, M., Mbua, E. and Nakatsukasa, M., 2015: New specimens of a primitive hippopotamus, *Kenyapotamus coryndonae*, from the Upper Miocene Nakali Formation, Kenya. *Journal of the Geological Society of Japan*, vol. 121, p. 153–159.
- van der Made, J., 1996: Listriodontinae (Suidae, Mammalia), their evolution, systematics and distribution in time and space. *Contributions to Tertiary and Quaternary Geology*, vol. 33, p. 3–254.
- von Meyer, H., 1846: Mitteilungen an Prof. Bronn gerichtet. *Neues Jahrbuch für Mineralogie, Geologie und Paläontologie*, vol. 1846, p. 462–476.
- Whitworth, T., 1958: Miocene ruminants of East Africa. *Fossil Mammals of Africa*, vol. 15, p. 1–50. British Museum (Natural History), London.

#### Figure and table captions

**Figure 1.** Map of Africa, showing the Nakali fossil locality in central Kenya (after Kunimatsu *et al.*, 2007). [planned for page width]

**Figure 2.** General stratigraphy and geological information of the Miocene Nakali Formation of central Kenya, showing the stratigraphic horizon of the specimens described here (after Kunimatsu *et al.*, 2007; Hilgen *et al.*, 2012). [planned for page width]

Figure 3. Cf. *Listriodon* sp. from the Miocene Nakali Formation of Kenya, KNM-NA 66161, a talonid (or distal lophid in the Suidae) fragment of a left lower molar. **A**, occlusal view (stereo pair); **B**, schematic drawing of occlusal view; **C**, lingual view; **D**, buccal view; **E**, ventral view; **F**, mesial view; **G**, distal view. [planned for page width]

Figure 4. *Dorcatherium* cf. *pigotti* Whitworth from the Miocene Nakali
Formation of Kenya, KNM-NA 57182, left DP4 or M1. A, occlusal view (stereo pair);
B, schematic drawing of occlusal view; C, lingual view; D, buccal view; E, mesial
view; F, distal view. [planned for page width]

**Figure 5.** *Dorcatherium* cf. *pigotti* Whitworth from the Miocene Nakali Formation of Kenya, KNM-NA 54202, a right mandible with p3–m3. **A**, lingual view; **B**, buccal view. [planned for page width] **Figure 6.** *Dorcatherium* cf. *pigotti* Whitworth from the Miocene Nakali Formation of Kenya, dentition (right p3–m3) of KNM-NA 54202. **A**, occlusal view (stereo pair) of right p3–m3; **B**, disto-occlusal view of right m3; B<sub>1</sub>, photo; B<sub>2</sub>, schematic drawing. Left scale corresponds to A, and right scale corresponds to B. [planned for page width]

**Figure 7.** *Dorcatherium* cf. *pigotti* Whitworth from the Miocene Nakali Formation of Kenya, p3–p4 of KNM-NA 54202. **A**, occlusal view; **B**, lingual view; **C**, buccal view. [planned for columnar width]

**Figure 8.** Upper and lower molars of a living tragulid species, *Tragulus javanicus* (Osbeck), in occlusal view. **A**, left M1–M2 of NMNS M31096 (male); **B**, right m2–m3 of NMNS M31096 with a hypoconulid basin on m3 (normal condition); **C**, right m3 of NMNS M31085 (male) without a hypoconulid basin (rare condition). [planned for columnar width]

**Table 1.** Dental measurements (in mm) of *Dorcatherium* cf. *pigotti* Whitworthfrom the Miocene Nakali Formation, Kenya. The p3 length of KNM-NA 54202 wasestimated based on the position of the mesial root. Abbreviations: *L*, mesiodistal length;*W*, buccolingual width; *W-m*, width of the mesial part; *W-d*, width of the distal part;*W-trd*, trigonid width; *W-tad*, talonid width.



















hypoconulid basin

5 mm



**Table 1.** Dental measurements (in mm) of *Dorcatherium* cf. *pigotti* Whitworth from the Miocene Nakali Formation, Kenya. The p3 length of KNM-NA 54202 was estimated based on the position of the mesial root. Abbreviations: *L*, mesiodistal length; *W*, buccolingual width; *W-m*, width of the mesial part; *W-d*, width of the distal part; *W-trd*, trigonid width; *W-tad*, talonid width.

Specimen	DP4 /M1	DP4 /M1	DP4 /M1	p3	р3	p4	p4	m1	m1	m1	m2	m2	m2	m3	m3	m3	m1–m3 row
KNM-NA	L	W-m	W-d	L	W	L	W	L	W-trd	W-tad	L	W-trd	W-tad	L	W-trd	W-tad	L
57182	7.5	7.1	7.0		_		_	_			_			_			
54202	_	_	_	> 7.4	3.5	7.2	3.7	7.9	4.7	5.0	8.8	5.7	6.2	12.8	6.4	6.6	29.6