

SHORT NOTES

An interesting mid-Cretaceous ammonite from Hokkaido (Studies of the Cretaceous ammonites from Hokkaido and Sakhalin-LXXIX)

TATSURO MATSUMOTO and NAOKO EGASHIRA

c/o Department of Earth and Planetary Sciences, Kyusyu University, Fukuoka, 812-81 Japan
1353 Yoga-cho, Saga, 840 Japan

Received 13 September 1996 ; Revised manuscript accepted 29 March 1997

Abstract. In this paper a fairly well-preserved ammonite is shortly described to show its interesting characters. It is provisionally referred to *Bhimaites* and regarded as a new species. Since it is small even at the adult stage and has a probable trace of lappet, it may represent a microconch of a dimorphic pair. It was found solely from the upper part of the Albian in the Soeushinai area of Hokkaido. More material is, however, required to clarify the extent of variation, to confirm the idea of dimorphism and to know the stratigraphic range and geographic distribution of this species.

Key words : Ammonite, *Bhimaites*, dimorphic pair, Hokkaido, mid-Cretaceous

Introduction

Among a large collection of ammonites from the mid-Cretaceous strata of the Soeushinai area, northwestern Hokkaido, there is a small but interesting ammonite which has not been reported previously. This is a short note on it.

Before going further, we greatly appreciate Yoshitaro Kawashita for his keen sight and also for the generous supply of his acquisition to scientific research. Thanks are extended to Tamio Nishida, University of Saga, for his warm help during the course of field and indoor works related to this study ; to C.W. Wright for his kindness in reading an earlier draft with linguistic improvement ; to Kazuko Mori for her assistance in preparing the manuscript.

Palaeontological description

Genus *Bhimaites* Matsumoto, 1954

Type species.—*Ammonites bhima* Stoliczka, 1865 (by original designation, Matsumoto, 1954, p. 113).

Remarks.—Because of the degree of morphological similarity, *Bhimaites* could be ranked among the subgenera of *Puzosia* Bayle, 1878. It contains several species of Albian and/or Cenomanian age. A few species have been recorded from the Turonian and possibly from the Coniacian. From the standpoint of phylogenetic divergence, the generic status of *Bhimaites* is kept at least for the time being (see Matsumoto, 1988, p. 21)

Species of *Bhimaites* are rare in the Cretaceous of Japan. *B. takahashii* Matsumoto (1988, p. 107, fig. 47) was a sole example from the mid-Turonian *Inoceramus hobetsensis* Zone in the Obira area (northwestern Hokkaido). Here is

another rare but interesting new species which is provisionally referred to *Bhimaites* with a query.

Bhimaites ? *kawai* sp. nov.

Figures 1-3

Material.—GK. H8490 (=YKC061111) from the laminated sandstone (feldspathic wacke) at locality R813 on the River Sounnai from the uppermost part of Member My2, Middle Yezo Group (see Nishida *et al.*, 1996, fig. 2 and fig. 10). This is the holotype. We have endeavored to get more material, but so far we have not been successful.

Diagnosis.—Shell small and discoidal, with moderate involution (about 3/5) and umbilical ratio (1/3). Whorls increasingly compressed with growth. Outer whorl narrowly oval in section with the maximum breadth at one third of height, from where the outer flanks converge gently to arched venter, while the inner flanks are bent abruptly to low but nearly vertical umbilical wall.

Phragmocone nearly smooth. Body chamber occupies about three-quarters of the last whorl. Constrictions fairly frequent on the body chamber, flexuous on the main part of the flank and much projected from the ventrolateral shoulder to the venter. On each interspace of the constrictions a few weak furrows may run roughly subparallel with the constrictions.

Suture rather simple ; L as deep as E and rather asymmetric ; U2 much smaller than L and tripartite ; auxiliaries still smaller and regularly descending.

Specific name.—This species is dedicated to Kawa-san [Mr. Kawa], a friendly nickname of Yoshitaro Kawashita, who found the holotype.




Figure 1. *Bhimaites ? kawai* sp. nov. Holotype. GK. H8490 [=YKC061111] from locality R813, Member My2 of the Soeushinai area. 1. left lateral (a), frontal (b), sectional (c), back (d) and right lateral views, $\times 1.7$. 2. left lateral view of the same specimen in different light before cleaning the umbilicus, $\times 1.5$; arrow: last septum. Bar scales: 10 mm for each of Figs. 1-1, 1-2.




Figure 2. *Bhimaites ? kawai* sp. nov. The last third suture (external) of the holotype. E, L, U2: external, lateral and 2nd umbilical lobes; s: umbilical seam. Bar scale: 1 mm.




Figure 3. *Bhimaites ? kawai* sp. nov. Cross-section (sketch) of the holotype. Bar scale: 2 mm. While the umbilicus was under cleaning, the specimen was cross-cut into halves and the section was sketched. The trace of the fissure may be seen in Figure 1-1a.

Dimensions.—See Table 1.

Observation.—The holotype is the sole material which is at present available for this species. Although its body chamber is fairly long, the actual apertural margin is not preserved. The trace of the faint furrow in front of the constriction shows a slightly more pronounced lateral convexity than that of the constriction. This seems to suggest the possible existence of a lappeted peristome. For the reason of the small size and the above observation, the holotype may represent a microconch. To confirm this presumption, examples of corresponding macroconchs should be searched out. It is noted that the holotype has a somewhat larger umbilical ratio (U/D) than that of the typical species of *Bhimaites*. This may be due to its small size, corresponding to a juvenile of a normal species of *Bhimaites*. Such a presumption should be examined by a find of macroconchs of this species.

Comparison and discussion.—In the previous literature, there is no species with which the present specimen can be

Table 1. Measurements of *Bhimaites ? kawai*.

Specimen	D	U	U/D	H	H/D	B	B/D	B/H	H/h
Holotype (E)	16.4	5.4	.33	6.8	.41	4.5	.28	.66	1.33
// (E-180°)	12.3	4.0	.33	5.1	.41	3.4	.28	.67	1.31

D=diameter, U=width of umbilicus, H=whorl-height, B=whorl-breadth, h=whorl-height at half a whorl adapical from H; E=preserved end, E-180°=at half a whorl adapical from E. Measurements of linear dimensions are in mm.

identified. The specimen from the presumed Lower Cenomanian of England, described as *Puzosia* (? *Bhimaites*) sp. by Wright and Kennedy (1984, p. 59, pl. 4, fig. 4; text-fig. 2) has compressed whorls and prorsiradiate constriction. It is, hence, somewhat similar to the Hokkaido specimen described above. It is still septate at its preserved end at a larger diameter (nearly 30 mm) than ours and is less compressed ($B/H=0.87$ as compared with 0.67 of our holotype) despite its later growth stage. The illustrated suture of the British specimen is rather of a *Puzosia* pattern, whereas that of ours is dissimilar to it, showing a somewhat peculiar feature.

The holotype of *Puzosia decemsulcata* Collignon (1964, p. 56, pl. 332, fig. 1495), from the "Lower Cenomanian" of Madagascar, has frequent constrictions which are prorsiradiate and slightly flexuous on the main part of the flank and markedly projected from the ventrolateral shoulder to the venter. In this respect it is fairly similar to our species, but it is larger (still septate at $D=90$ mm) and has a larger ratio B/H [0.78] than that of our species. Collignon's species may be referable to *Bhimaites*, but a final conclusion should be deferred, until it is defined more clearly.

As in the case of *Microdesmoceras tetragonum* Matsumoto and Muramoto in the Desmocerotinae (see Matsumoto *et al.*, 1972), this species might represent a progenetic dwarf offshoot of *Puzosia*, but we hesitate to propose a new subgenus for the reason of insufficient evidence.

Occurrence.—As for *Material*. Member My2 is referred to the upper part of the Upper Albian (see Nishida *et al.*, 1996). The true stratigraphic range and geographic distribution of this species should be determined by further investigations.

References cited

- Collignon, M., 1964: *Atlas des fossiles caractéristiques de Madagascar (Ammonite)*. Fascicle 11 (Cenomanien). Service Géologique, Tanannarive. p. i-xi, 1-152, pls. 318-375.
- Matsumoto, T., 1954: Family Puzosiidae from Hokkaido and Saghalien. *Memoirs of the Faculty of Science, Kyushu University, Series D*, vol. 5, no. 2, p. 69-118, pls. 9-23.
- Matsumoto, T. (compiler), 1988: A monograph of the Puzosiidae (Ammonoidea) from the Cretaceous of Hokkaido. *Palaeontological Society of Japan, Special Papers*, no. 30, p. i-iii, 1-179, 88 figs.
- Matsumoto, T., Muramoto, T. and Inoma, A., 1972: Two small desmocerotid ammonites from Hokkaido. *Transactions and Proceedings of the Palaeontological Society of Japan, New Series*, no. 87, p. 377-394, pl. 47.
- Nishida, T., Matsumoto, T., Yokoi, K., Kawashita, Y., Kyuma, Y., Egashira, N., Aizawa, J., Maiya, S., Ikuji, Y. and Yao, A., 1996: Biostratigraphy of the Cretaceous Middle Yezo Group in the Soeushinai area of Hokkaido— with special reference to the transitional part from Lower to Upper Cretaceous. *Journal of Faculty of Education, Saga University*, vol. 43, no. 1, p. 63-147. (*in Japanese with English abstract*)
- Stoliczka, F., 1865: The fossil Cephalopoda of the Cretaceous rock of southern India. Ammonoidea, with revision of the Nautilidae. *Memoirs of the Geological Survey of India, Palaeontologia Indica, series 3*, p. 107-154, pls. 55-80.
- Wright, C.W. and Kennedy, W.J., 1984: The Ammonoidea of the Lower Chalk. Part 1. *Monograph of the Palaeontological Society*, London, p. 1-126, pls. 1-4.

A GUIDE FOR PREPARING MANUSCRIPTS

PALEONTOLOGICAL RESEARCH is dedicated to serving the international community through the dissemination of knowledge in all areas of paleontological research. The journal publishes articles, normally not exceeding 24 pages, and short notes, normally less than 4 pages. Manuscripts submitted are subject to review and editing by reviewers and a language editor. A condition of acceptance for all manuscripts is that they have not been published and will not be submitted nor published elsewhere. Manuscripts accepted for publication will generally be published in order of submission. Authors submit three copies of their manuscript for editorial review. After review, two copies of the revised manuscript are to be returned for copy editing.

Text : Paleontological Research is intended to be read by an international audience, therefore it is particularly critical that language be clear and concise. Though manuscripts in French and German may occasionally be accepted (with abstract and captions for illustrations and tables also provided in English), English is strongly favored as the language of international scientific communication. For manuscripts prepared in English, either British or American usage in style, words and spelling is acceptable. Use SI (Système International d'Unités) units wherever possible.

Text should be typed always in double space on one side of opaque white paper of not less than either 210×280 mm (A4 size) or 8 1/2×11 inches in the following order.

Cover sheet. Cover sheet should contain (1) full name, address, phone number and fax number of the author taking responsibility for the galley proofs, (2) running title composed of less than 40 characters, and (3) the numbers of tables and figures.

Title page. Title of the paper, names of authors and their professional affiliations with addresses (or residential address, if an author is unaffiliated). Titles are to be brief and simple. Spell out one or more of the authors' first names.

Abstract. Abstract should be a condensation and concentration of the essential qualities of the paper. All the papers, including Short Notes, are to be accompanied by an abstract not exceeding 500 words. New taxonomic or stratigraphic names should be mentioned in the abstract.

Key words. Select keywords (not more than six words or phrases) which identify the most important subjects covered by the paper and arrange them in alphabetical order.

Main text. Use three or fewer levels of heading. No footnotes are to be used. Bibliographical references are to be identified by citing the authors' names, followed, in parentheses, by the date of publication, with a page number if

desired. All citations must have a corresponding entry in the reference list.

The typical format for arrangement of systematic paleontology can be learned from current issues of the Journal. All descriptions of new taxa must include a diagnosis, and, as appropriate, stratigraphic and geographic indications, designation of a type or types, depository information, and specification of illustrations. In synonymies use an abbreviated form of the reference, consisting only of authors of reference, date of publication, and number of pages, plates, figures and text-figures referring to the organism or organisms in question.

References. Heading for the bibliography can be either "References cited" or "References." Entries are to be listed alphabetically. No abbreviations will be used in article and book titles. Journal titles are written out, not abbreviated. Series, volume, and number or part are to be given, with the appropriate word abbreviated in each case ("ser.", "vol.", etc.; see the next page for examples).

Illustrations : All illustrations, including maps, geologic sections, and half-tone illustrations (including "plates") are to be called figures and must be numbered in the same sequence as they are first cited in the text. Citations of illustrations in the text are to be spelled out in full (e.g., Figure 2 or Figure 2.1). Figure captions are to be typed separately. Plan the illustrations so that they take up either the entire width of the printed page (170 mm) or the width of one column (80 mm). Originals should not be smaller than the final intended size for printing. No foldouts will be accepted. Mark all originals clearly with authors' names and figure number. Photographs of all specimens except sections must be illuminated from the upper left side, as is conventional.

Manuscripts on disk : Authors are encouraged to deliver final, revised manuscript copy on disk, but disks should be sent only after the paper has been accepted. Both 5.25 and 3.5 inch disks with the text in a recent version of Word Perfect or Microsoft Word are acceptable. Be sure to specify, in a covering note, the hardware and the word-processing package used.

Galley proofs and offprints : Galley proofs will be sent to authors about one month before the expected publication date and should be returned to the Editors within 3 days of receipt. The authors are responsible for reading the first galley proof. Minor changes submitted by the author will be permitted while a paper is in galleys, but a charge will be made for substantial alterations.

The authors receive free of charge 50 offprints without covers. Additional copies and covers can be purchased and should be ordered when the proofs are returned.

Charges : If a paper exceeds 24 printed pages, payment of page charges for the extra pages is a prerequisite for acceptance. Illustrations in color can also be published at the authors' expense. For either case, the Editors will provide information about current page charges.

Return of published figures : The manuscripts of the papers published will not be returned to the authors. However, figures will be returned upon request by the authors after the paper has been published.

- Ager, D.V., 1963: *Principles of Paleocology*, 371 p. McGraw-Hill Co., New York.
- Barron, J.A., 1983: Latest Oligocene through early Middle Miocene diatom biostratigraphy of the eastern tropical Pacific. *Marine Micropaleontology*, vol. 7, p. 487-515.
- Barron, J.A., 1989: Lower Miocene to Quaternary diatom biostratigraphy of Leg 57, off northeastern Japan, Deep Sea Drilling Project. In, Scientific Party, *Initial Reports*

- of the Deep Sea Drilling Project*, vols. 56 and 57, p. 641-685. U.S. Govt. Printing Office, Washington, D.C.
- Burckle, L.H., 1978: Marine diatoms. In, Haq, B.U. and Boersma, A. eds., *Introduction to Marine Micropaleontology*, p. 245-266. Elsevier, New York.
- Fenner, J. and Mikkelsen, N., 1990: Eocene-Oligocene diatoms in the western Indian Ocean: Taxonomy, stratigraphy, and paleoecology. In, Duncan, R.A., Backman, J., Peterson, L.C., et al., *Proceedings of the Ocean Drilling Program, Scientific Results*, vol. 115, p. 433-463. College Station, TX (Ocean Drilling Program).
- Koizumi, I., 1972: Marine diatom flora of the Pliocene Tatsunokuchi Formation in Fukushima Prefecture. *Transactions and Proceedings of the Palaeontological Society of Japan, New Series*, no. 86, p. 340-359.
- Kuramoto, S., 1996: Geophysical investigation for methane hydrates and the significance of BSR. *Journal of the Geological Society of Japan*, vol. 11, p. 951-958. (in Japanese with English abstract)

Palaeontological Society of Japan (JSP) Standing Committee Actions

During its meeting on March 8, the JSP Standing Committee enacted the following changes to its membership.

- New members elected ;
 Tuyahiko Fujimoto, Hiroaki Yoshikawa.
- Resigned members ;
 Masayoshi Asaga, Hidehiko Asai, Yukimasa Oho,
 Akira Okajima, Naomi Oyama, Kazuhisa Satomi,
 Sadako Takeuti.
- Deceased member ;
 Sotoji Imamura.

During its meeting on May 24, the JSP Standing Committee enacted the following changes to its membership.

- New members elected ;
 Kohsaku Arai, Yoshimasa Date, Shinjiro Fujimoto,
 Kazuhiko Fujita, Takao Inoue, Toshiyuki Kurihara,
 Jianbo Liu, Yuriko Matsumura, Takao Mukuda,
 Hiroyoshi Sano, Masahiro Shiraki, Yuji Soeda,
 Hiroaki Tanaka, Bunji Tojo, Junichi Uchida,
 Atsuko Uchino.
- Resigned members ;
 Ikuo Fujita, Noriyuki Nishida, Morihiko Ofusa,
 Japan Energy Development CO., LTD.